MAV Mathematical Methods Trial CAT 3 1999

Question 1

Three brothers Adam, Mark and Paul Smith inherit a family farm. The farm has four trees on its boundary. The brothers decide to use the trees to divide the land into three separate farms. The northernmost tree is a large gum tree. The other three trees are a wattle, a cedar and a fig. the proposed d These three tre

M is defined as A is defined as

P is defined as Distances are a

2 marks	

2 m		

© Mathematical Ass	© Mathematical Association of Victoria 1999 Page 1
	2 Hains
	2 montes
	s. What is the equation of the boundary line mixing the cenal tree and the goin tree:
	2 marks
	THE DOME HE COMMITTEE OF THE COMMITTEE AND BOTH HAVE
	Write down the coordinates of the redar tree and the gum tree
	Distances are all measured in kilometres.
	s is defined as the area inherited by Paul.
	If is defined as the area inherited by Mark.
ring ine	Count Fig.
expresse	Cata
e. The equi	$O \downarrow 1$ 2×3 4 5 6 7×8 $East(x)$
	2
	3 👯
	4 Coum
)
	6.1
d. Which I	7
	North(t) 8 ♣
	0, 0) as shown below.
	he farm can be represented by a set of cartesian axes with the Wattle situated at the origin.
	he proposed division of the land.
	hese three trees all stand on an East-West line. The Wattle is taken as the reference point for

NIAV Mathematical Methods Trial CAT 3 1999 What is the distance between the cedar tree and the gum tree? uation of the boundary line linking the wattle, gum and fig trees can be ed as a quadratic equation, $y = ax^2 + bx + c$, where a, b and c are constants c exact values for a, b and c. ree is further away from the cedar tree, the gum tree or the fig tree? 4 marks 1 mark l mark

© Mathematical Association of Victoria 1999

Page 2

[working space]	4 marks Total 14 Marks			f. Using the method of integration, find the area of Mark's farm correct to two decimal places.

MAV Mathematical Methods Trial CAT 3 1999

Question 2

A new ferris wheel at a theme park revolves at a constant speed in an anti-clockwise direction, requiring passengers to enter and exit the wheel as it is in motion. For safety, passengers enter the ferris wheel at its lowest point.

The height, in metres, of a passenger above the ground t minutes after they have entered the ferris wheel is given by

$$(t) = 12 - 10\cos\frac{\pi}{2}t, \quad t \ge$$

	20
whole minutes does a ride take?	A "ride" on this ferris wheel finishes after two revolutions of the wheel. How many
	<u>ټ</u>

b. What	
What is the maximum height passengers can reach on the wheel?	
eight passengers can	
1 reach on the whee	
2 mark	

		r
	that passengers enter the wheel without stepping up or down?	The passengers enter the wheel from a platform. What height must this platform be so

l mark


	,	
l mark		

- 1	,
	£
- 1	7
- 1	
- 1	Ĭ
- 1	₹
- 1	į
- 1	3
- 1	캽
- 1	5
- 1	÷
- !	÷
- [2
- 1	•
-1	₹
ſ	Ξ
ŀ	O
- }	_
- [2
- 1	_
- 1	\$
- 1	•

d. Show that it takes 1.41 minutes for passengers on the wheel to reach a height of 18 metres for the first time?

l mark

The ferris wheel and its surroundings can be represented on a set of cartesian axes as shown below. In this diagram, y is defined as the height above ground and x is the horizontal distance from the centre of the platform. The distances are measured in metres.



e. Show that the equation of the line joining the point A on the ground and the point B on the edge of the top of a group of buildings can be given by

$$y=\frac{1}{7}(134-x)$$

2 marks		i	

© Mathematical Association of Victoria 1999

MAV Mathematical Methods Trial CAT 3 1999

The position of a passenger can be given by the relation

$$(y-12)^2 + x^2 = 10^2$$

Show that the points of intersection between this relation and the line in part f are (-6, 20) and (8, 18).

g. For what percentage of time on the ferris wheel will a passenger be able to see the point A? Assume that a passenger's line of sight begins from a point on the ferris wheel making no allowance for how tall the passenger is. Write your answer to the nearest whole percentage.

3 marks

Total 13 marks

l	Ā
	į
l	Metho
Ì	ds Tri
ı	EL CA
l	3
	999

Question 3

The height, h(x), of a water slide above the ground is given by

$$h_1(x) = 3 - \frac{1}{4}x^2$$

 $h_2(x) = Ae^{-k(x-1)} - 0.5,$

 $1 < x \le 7$, where x is the horizontal distance in metres from the entrance point at the top of the slide and A and k are real constants

Given that the functions h_1 and h_2 intersect at x = 1, show that A = 3.25.

2 marks

Find $h_1'(x)$, the derivative of $h_1(x)$.

Ģ

l mark

;=: Find $h_2'(x)$, the derivative of $h_2(x)$.

l mark

፷፧ The water slide is continuous and smooth at the intersection between the functions, h_1 and h_2 . Use this information to find the constant k correct to two decimal places.

2 marks

Page 7

© Mathematical Association of Victoria 1999

MAV Mathematical Methods Trial CAT 3 1999

At what height, h, would a person be if they slide along the slide for a horizontal distance of 2 metres? Write your answer correct to two decimal places.

What horizontal distance would a person travel after having descended a height of 2 2 marks

metres? Write your answer accurate to the nearest second decimal place

ġ.

ç Draw a graph of the functions, h_1 and h_2 on the following set of axes.

2 marks

0 *4* √ N 6 2 marks

© Mathematical Association of Victoria 1999

Page 8

4 marks		4 marks	Total 16 marks	[working space]		
a. What is the probability i. more than 65 n ii. less than 65.8 r b. Write an expression in probability of throwing				c. Use the expression fro		

1 mark												
												I
Write	greater.	tres or	65 me	throws	ber of t	ne num	ı, for tl	portion three d	ple pro	he sam	Find \hat{p} , the sample proportion, for the number of throws 65 metres or greater. Write your answer correct to three decimal places	~ _
-	0		22	2/	26	39	2/	22	=	· ·	Number of Throws	₹3
68	67	1 8	3 8	42	88	62	62	8	. 59	88	Distance	1 <u>2</u>
and own in	e games tre as sh	fore the	eeks be ne near	ts to th	s injure 1e resul	ally wa	that S. Ie reco	spected rows. F	ies, sus	bert Jor 200 of le.	Sally's coach, Albert Jones, suspected that Sally was injured 2 weeks before the games and secretly recorded 200 of her throws. He recorded the results to the nearest metre as shown in the following table.	llo s c
3 marks											i	
												1 1
of	distance in.	or the . Expla	terval f record?	ility int a new r	probab create a	e 95% ely to	ace, the	imal pl think sl	ne dec o you t	ect to c	Find, correct to one decimal place, the 95% probability interval for the distance of Sally's throws. Do you think she is likely to create a new record? Explain.	IO TI
3 marks												
												İ
									!			
												İ
final?	nake the	y will n	at Sally	ces, tha	mal pla	∕o deci	e to tw	ассига	bility,	e proba	What is the probability, accurate to two decimal places, that Sally will make the final?	
rd is	The reco	etres. 1	d 63 m	arounc	d to be	xpecte	nal is e	mpic fi	hé Olyi	rk for t	The qualifying mark for the Olympic final is expected to be around 63 metres. The record is 72.3 metres.	[[
								3	CAT 3 19	rods Trial	tAV Mathematical Methods Trial CAT 3 1999	- 1 →

Find, correct to three decimal places; the 95% confidence proportion, p, for the final two weeks of Sally's perform. With reference to p or pexplain whether Albert's suspi		F	úé
of p.	With reference to p or $\stackrel{\wedge}{p}$ explain whether Albert's suspicions are true.	l m Find, correct to three decimal places; the 95% confidence interval for the population proportion, p, for the final two weeks of Sally's performance.	Find, correct to three decimal places, the standard error of \hat{p} .

END OF TRIAL CAT 3

ociation of Victoria 1999