

SECTION 1

1	2	3	4	5	6	7	8	9	10	11
C	E	A	C	D	B	D	A	B	A	E

12	13	14	15	16	17	18	19	20	21	22
D	E	D	B	A	C	B	A	C	C	D

Q1 $\tan \theta = \frac{x}{x-1}$, $\tan \phi = 1 - \frac{1}{x} = \frac{x-1}{x}$, $\therefore \tan \phi = \frac{1}{\tan \theta}$

$\therefore \theta + \phi = \frac{\pi}{2}$ C

Q2 $f(x) = e^{\log_e \frac{1}{e} \sqrt{x}} = e^{\frac{\log_e \sqrt{x}}{\log_e \frac{1}{e}}} = e^{-\log_e \sqrt{x}} = e^{\log_e x^{-\frac{1}{2}}} = x^{-\frac{1}{2}}$ E

Q3 $x=0$, $nx=\pm\pi$, $nx=\pm\frac{\pi}{2}$

$\therefore x=0$, $x=\pm\frac{\pi}{n}$, $x=\pm\frac{\pi}{2n}$

Q4 $\int_2^{10} f(x)dx = 10 \times 10 - 28 - 4 \times 2 = 64$

Q5 x-intercepts: $ax^2 - 1 = 0$, $x = \pm\frac{1}{\sqrt{a}}$

Area = $-\int_{-\frac{1}{\sqrt{a}}}^{\frac{1}{\sqrt{a}}} (ax^2 - 1)dx = \int_{-\frac{1}{\sqrt{a}}}^{\frac{1}{\sqrt{a}}} (1 - ax^2)dx$ D

Q6 $x^2 - a^2 \geq 0$, $a^2 - x^2 \geq 0$ and $x+a \neq 0$, $\therefore x=a$

Q7

A Q14 $\int f(x)dx = 1 - 2x - \frac{1}{4} \log_e(1-2x)$

$f(x) = \frac{d}{dx} \left(1 - 2x - \frac{1}{4} \log_e(1-2x) \right) = -2 + \frac{1}{2} \times \frac{1}{1-2x}$ D

Q15 $\int_0^{\frac{\pi}{8}} g(x)dx = \int_0^{\frac{\pi}{8}} \frac{2}{\cos^2(2x)} dx = \int_0^{\frac{\pi}{8}} 2 \sec^2(2x)dx$
 $= [\tan(2x)]_0^{\frac{\pi}{8}} = \tan \frac{\pi}{4} - \tan 0 = 1$


Average value = $\frac{\int_0^{\frac{\pi}{8}} g(x)dx}{\frac{\pi}{8}-0} = \frac{1}{\frac{\pi}{8}} = \frac{8}{\pi}$ B

Q16 f is a many-to-one function, $\therefore f^{-1}$ does not exist. A

A Q17 The last draw is equally likely to be blue, green or red,
 $\therefore \Pr(\text{last.draw.is.red}) = \frac{1}{3}$ C


Q18

Q19



$\Pr(\text{Home.between.8.and.9})$
 $= \Pr(\text{left.between.6.and.9}) - \Pr(\text{left.between.6.and.8})$
 $= 1 - \frac{7}{8} = \frac{1}{8} = 0.125$ A

Q10



Q20

Transition matrix $\begin{bmatrix} 0.85 & 0.10 \\ 0.15 & 0.90 \end{bmatrix}$, state matrix $\begin{bmatrix} 0.65 \\ 0.35 \end{bmatrix}$.

State matrix after 2 weeks: $\begin{bmatrix} 0.85 & 0.10 \\ 0.15 & 0.90 \end{bmatrix}^2 \begin{bmatrix} 0.65 \\ 0.35 \end{bmatrix} = \begin{bmatrix} 0.54 \\ 0.46 \end{bmatrix}$

C

Q21 $\sum p(x) = 1$

$\therefore p(0.01) + p(0.10) + p(0.30) + p(0.50) + p(0.59) = 1$

$\therefore 5a + 1.009 = 1, \therefore a = -0.0018$

$$\bar{X} = \sum xp(x)$$

$= 0.01p(0.01) + 0.10p(0.10) + 0.30p(0.30) + 0.50p(0.50) + 0.59p(0.59)$

≈ 0.30

C

Q22 $\mu = 12.3, \sigma = \sqrt{1.69} = 1.3$

$\Pr(9.1 \leq X < b) = \Pr(X < b) - \Pr(X < 9.1) = 0.95$

$\Pr(X < b) = 0.95 + 0.006917 \approx 0.9569, \therefore b \approx 14.53$

D

SECTION 2

Q1a $y = f(x) = e^x - 2x$

Let $f'(x) = e^x - 2 = 0, \therefore e^x = 2, x = \log_e 2$

$\therefore y = 2 - 2\log_e 2 = 2(1 - \log_e 2)$

f has a stationary point, $(\log_e 2, 2(1 - \log_e 2))$.

Q1b

x	0	$\log_e 2$	1
$f'(x)$	negative	zero	positive

$(\log_e 2, 2(1 - \log_e 2))$ is a minimum.

Q1c Since $(\log_e 2, 2(1 - \log_e 2))$ is a minimum and

$2(1 - \log_e 2) > 0, \therefore e^x - 2x > 0$ for $x \in R$

$\therefore e^x > 2x$ for $x \in R$.

Q1d When $x = 0, e^x = 1$ and $x^2 + 1 = 1$

$\therefore e^x = x^2 + 1$ at $x = 0$, i.e. both functions have a common point at $x = 0$.

$\frac{d}{dx} e^x = e^x$ and $\frac{d}{dx}(x^2 + 1) = 2x$

Since $e^x > 2x$ for $x \in R, \therefore e^x > 2x$ for $x > 0$, i.e. the rate of increase of e^x is greater than the rate of increase of $x^2 + 1$.

$\therefore e^x \geq x^2 + 1$ for $x \geq 0$

Q1e $(x-1)^2 \geq 0, x^2 - 2x + 1 \geq 0, \therefore x^2 + 1 \geq 2x$

Q1f $\frac{d}{dx} \log_e(x^2 + 1) = \frac{d}{du} \log_e(u) \times \frac{d}{dx}(x^2 + 1) = \frac{2x}{x^2 + 1}$

Q1h $\int_0^x \frac{2t}{t^2 + 1} dt = [\log_e(t^2 + 1)]_0^x = \log_e(x^2 + 1)$

Q1i From part e, $t^2 + 1 \geq 2t, \therefore 1 \geq \frac{2t}{t^2 + 1}$

$\therefore \int_0^x 1 dt \geq \int_0^x \frac{2t}{t^2 + 1} dt$ for $x \geq 0$

$\therefore [t]_0^x \geq \log_e(x^2 + 1), \therefore x \geq \log_e(x^2 + 1)$

Hence $e^x \geq x^2 + 1$ for $x \geq 0$

Q2a $P(x) = x^3 + 6ax^2 + 6bx + 4c$

$P'(x) = 3x^2 + 12ax + 6b$

Q2b The turning point is on the x -axis, $\therefore P'(x) = 0$ and $P(x) = 0$

$P'(x) = 0 \therefore 3x^2 + 12ax + 6b = 0, x^2 + 4ax + 2b = 0,$

$\therefore x^3 + 4ax^2 + 2bx = 0$ for $x \neq 0 \dots \dots \dots (1)$

$ax^2 + 4a^2x + 2ab = 0$ for $a \neq 0 \dots \dots \dots (2)$

$P(x) = 0, \therefore x^3 + 6ax^2 + 6bx + 4c = 0 \dots \dots \dots (3)$

$(3) - (1): 2ax^2 + 4bx + 4c = 0, \therefore ax^2 + 2bx + 2c = 0 \dots \dots \dots (4)$

$(2) - (4): 4a^2x - 2bx + 2ab - 2c = 0$

$\therefore (2a^2 - b)x + ab - c = 0$

$\therefore x = \frac{c - ab}{2a^2 - b}$

\therefore the turning point on the x -axis is $\left(\frac{c - ab}{2a^2 - b}, 0\right)$.

Q2ci Compare $Q(x) = x^3 + 0.4x^2 - 3.36x - 2.88$ with

$P(x) = x^3 + 6ax^2 + 6bx + 4c$

$a = \frac{1}{15}, b = -0.56$ and $c = -0.72$

$\therefore x = \frac{c - ab}{2a^2 - b} = -1.2$

\therefore the turning point on the x -axis is $(-1.2, 0)$.

Q2cii $Q(x) = x^3 + 0.4x^2 - 3.36x - 2.88 = (x + 1.2)^2(x - p)$

$\therefore 1.2^2 p = 2.88, p = 2$

\therefore the x -coordinate of the other x -intercept is 2

$Q'(x) = 3x^2 + 0.8x - 3.36 = 3(x + 1.2)(x - q) = 0$

$\therefore 3.6q = 3.36, q = \frac{14}{15}$


\therefore the x -coordinate of the other turning point is $\frac{14}{15}$.

Q2d Area $= - \int_{-1.2}^2 Q(x) dx \approx 8.74$ by CAS/graphics calc.

Q3a $y = a \log_e x + b$
 $(1,0)$, $0 = a \log_e 1 + b$, $b = 0$
 $(e,1)$, $1 = a \log_e e + 0$, $a = 1$
 \therefore curve B has the equation $y = \log_e x$.

Q3b $x^2 + (y-1)^2 = 1$, where $x \in [-1,1]$ and $y \in [0,1]$
 $(y-1)^2 = 1 - x^2$, $y-1 = -\sqrt{1-x^2}$
 \therefore curve A has the equation $y = 1 - \sqrt{1-x^2}$.

Q3c



Q3d Area of the ground space

$$= \text{rectangle } (e \times 1) - \text{quarter circle } \left(\frac{\pi \times 1^2}{4} \right) - \int_1^e \log_e x dx \\ = e - \frac{\pi}{4} - 1$$

Q3ei Curve B : $y = \log_e x$, at $x = p$, $y = \log_e p$

$$\frac{dy}{dx} = \frac{1}{x}, \text{ at } x = p, \frac{dy}{dx} = \frac{1}{p}, \therefore \text{gradient of the normal} = -p$$

Equation of the normal at $x = p$:

$$y - \log_e p = -p(x - p)$$

$$y = -px + p^2 + \log_e p$$

Q3eii $y = -px + p^2 + \log_e p$

Centre $(0,1)$, $1 = p^2 + \log_e p$, $\therefore p = 1$

\therefore equation of the normal at $x = 1$ is $y = -x + 1$.

Q3eiii The normal is perpendicular to both curves, \therefore the distance between the two curves along the normal is the shortest.
Shortest distance = distance between $(0,1)$ and $(1,0)$ – radius
 $= \sqrt{(1-0)^2 + (0-1)^2} - 1 = \sqrt{2} - 1$

Q3fi Curve B : $\frac{dy}{dx} = \frac{1}{x}$, curve A : $\frac{dy}{dx} = \frac{x}{\sqrt{1-x^2}}$

At $y = c$, $x = e^c$ for curve B , and $x = \sqrt{1-(c-1)^2}$ for curve A .

$$\therefore \frac{1}{e^c} = \frac{\sqrt{1-(c-1)^2}}{\sqrt{(c-1)^2}}, c = 0.22007 \approx 0.22 \text{ by CAS/graphics calc.}$$

Q3fii At $y = c = 0.22007$,
 $x = e^{0.22007} \approx 0.62587$ for curve A ,
and $x = \sqrt{1 - (0.22007 - 1)^2} \approx 1.24616$ for curve B .
Magnitude of translation = $1.24616 - 0.62587 \approx 0.62$

Q4ai $\Pr(\mu - w < X < \mu + w) = 0.8000$,
 $\therefore \Pr(X < \mu - w) = \frac{1 - 0.8000}{2} = 0.1000$, $\mu - w = 1.9468$
 $w = \mu - 1.9468 = 1.9500 - 1.9468 = 0.0032$

$$Q4aii (1 - 0.8000) \times 1000 = 200$$

$$Q4aiii \Pr(\mu - 2\sigma < X < \mu + 2\sigma | X < \mu - w \cup X > \mu + w) \\ = \frac{\Pr((\mu - 2\sigma < X < \mu + 2\sigma) \cap (X < \mu - w \cup X > \mu + w))}{\Pr(X < \mu - w \cup X > \mu + w)} \\ = \frac{0.9545 - 0.8000}{1 - 0.8000} \approx 0.7725$$

$$Q4bi \mu = 1.9500 \times 4 = 7.8000 \\ \sigma^2 = 0.0250^2 \times 4 = 0.0025, \therefore \sigma = 0.0500$$

$$Q4bii \Pr(X > 7.8750) \approx 0.0668$$

Q4biii Binomial distribution: $n = 26$, $p = 0.0668$
 $\Pr(X > 2) = 1 - \Pr(X \leq 2) \approx 1 - 0.7500 = 0.2500$

Q4biv No extra postage is required if the letter is $7.8750 - 0.1500 = 7.7250$ grams or less.
 $\Pr(X < 7.7250) = 0.0668$

$$Q4ci \int_{1.5000}^{2.2000} ke^{\frac{x}{2}} \sin(x) dx = 1, \int_{1.5000}^{2.2000} e^{\frac{x}{2}} \sin(x) dx = \frac{1}{k}, \\ k = 0.602036 \approx 0.6020$$

$$Q4cii \mu = \int_{1.5000}^{2.2000} x \times 0.602036 e^{\frac{x}{2}} \sin(x) dx = 1.8582344 \approx 1.8582$$

Q4ciii Let m grams be the median weight.

$$\int_{1.5000}^m 0.602036 e^{\frac{x}{2}} \sin(x) dx = 0.5, m = 1.86224 \approx 1.8622 \text{ by CAS/graphics calc.}$$

$$Q4civ \Pr(X > m | X > \mu) = \frac{\Pr(X > m \cap X > \mu)}{\Pr(X > \mu)} \\ = \frac{\Pr(X > m)}{\Pr(X > \mu)} = \frac{0.5}{\int_{1.8582344}^{2.2000} 0.602036 e^{\frac{x}{2}} \sin(x) dx} \approx 0.9884$$

Please inform mathline@itute.com re conceptual,
mathematical and/or typing errors