

MATHEMATICAL METHODS (CAS)

Units 3 & 4 – Written examination 2



2011 Trial Examination

SOLUTIONS

SECTION 1: Multiple-choice questions (1 mark each)

Question 1

Answer: E

Explanation:

$$\text{If } k = 2: (2 - 1)x + 4y = 8 \text{ and } 3x - (-2 + 2)y = 2 + 1$$

$$x + 4y = 8 \text{ and } 3x = 3$$

$$x = 1 \text{ and } y = \frac{7}{4}$$

Therefore it will have a unique solution.

Question 2

Answer: C

Explanation:

$$f(x - y) = (x - y)^3$$


$$f(x) - f(y) = x^3 - y^3$$

$$f(x - y) \neq f(x) - f(y)$$

Question 3

Answer: B

Explanation:
Sketch graph:



Question 4

Answer: E

Explanation:

Rewrite the equations and then use matrix form.

$$2x + 0y + 2z = -1$$

$$0x - 2y + 2z = 0$$

$$2x + 2y + 0z = -4$$

Question 5

Answer: B

Explanation:

$$m_{\text{tangent}} = -3$$

$$y = -3(x - 3) + 4 - 6 = -3x + 7$$

Question 6

Answer: B

Explanation:

$$Pr(Z < -z) = Pr(Z > z) = 0.75$$

2011 MATHMETH (CAS) EXAM 2

Use CAS Probability menu: inverse normal. Change the area to 0.25 because the TI-nspire calculators use $Z < c$ and not $Z > c$.

$$c = -0.6745$$

Question 7

Answer: B

Explanation:

There is a gap at $(0, -3)$ not a cusp.

Question 8

Answer: A

Explanation:

$f \circ g$ exists if $\text{ran } g \subseteq \text{dom } f$, therefore $D = x < -1$ or $x > 3$.

(For $f \circ g$ to exist, the range of g : $(1, \infty)$)

Question 9

Answer: C

Explanation:

Use CAS: $f(x) = \int (2-x)(2x+1)^2 dx = -x^4 + \frac{4}{3}x^3 + \frac{7}{2}x^2 + 2x$

Question 10

Answer: B

Explanation:

Solve $\int_k^{\frac{\pi}{2}} \frac{2}{\pi} \cos^2(x) dx = 0.3$ on CAS

$$k = 2.63$$

Question 11

Answer: B

Explanation:

$$\int_0^6 f\left(\frac{1}{3}x\right) + 1 dx = \int_0^6 f\left(\frac{x}{3}\right) dx + \int_0^6 1 dx = 12 + [x]_0^6 = 12 + 6 = 18$$

Question 12

Answer: C

Explanation:

$$\Pr(-1 < Z < 2) = \Pr(\mu - \sigma < X < \mu + 2\sigma).$$

Question 13

Answer: D

Explanation:

Use CAS: $\frac{d}{dx}(e^{-4x} \sin(x - 2))|_{x=2}$

Question 14

Answer: C

Explanation:


Use CAS and solve $\cos(2x) = -\frac{\sqrt{3}}{2}$, then expand to get $\pm \frac{5\pi}{12} + k\pi, k \in Z$

Question 15

Answer: E

Explanation:

Use CAS to sketch graph and find local maximum, then $(-6, -1.5) \cup (3, \infty)$.



Question 16

Answer: E

Explanation:

$$y = -5e^{2(x+2)} + 2$$

$$T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 5 \end{bmatrix} + \begin{bmatrix} -2 \\ 2 \end{bmatrix}$$

Question 17

Answer: B

Explanation:

Use CAS: Binomial pdf (20, 0.45, 5), $\Pr(X = 5) = 0.0365$

Question 18

Answer: A

Explanation:

$$\text{Period} = \frac{\pi}{b} = 5\pi, \text{ asymptote } x = \frac{\pi}{2b} = \frac{5\pi}{2}$$

Question 19

Answer: C

Explanation:


$$l_1 = 3 - e^{1-1} = 3 - 1 = 2, \quad l_2 = 3 - e^{2-1} = 3 - e$$
$$A = 2 + 3 - e = 5 - e$$

Question 20

Answer: A

Explanation:

Use CAS to sketch the graph.



There is a vertical line at $x = 0$.

Question 21

Answer: C

Explanation:

You have to use the points of intersection and for each section it is the top graph minus the bottom graph.

Question 22

Answer: E

Explanation:

The graph has a vertical asymptote at $x = a$ and the function lies to the right of the asymptote, sketch the graph.

SECTION 2: Analysis Questions

Question 1

a. $f(x) = \frac{2}{x-3} - 4 = 2(x-3)^{-1} - 4$

Use chain rule: $m_{\text{tangent}} = f'(x) = -2(x-3)^{-2} \times 1 = \frac{-2}{(x-3)^2}$

$m_{\text{normal}} = \frac{(x-3)^2}{2}$

M1+A1
2 marks

b. $m = 2, (1, -5)$

$y - y_1 = m(x - x_1)$

$y + 5 = 2(x - 1)$


$g(x) = 2x - 7$

M1+A1
2 marks

c. $g(f(x)) = g\left(\frac{2}{x-3} - 4\right) = 2\left(\frac{2}{x-3} - 4\right) - 7 = \frac{4}{x-3} - 15$, the domain is the set of all values in the domain of f (the domain of the inner function). Domain: $x \in \mathbb{R} \setminus \{3\}$.
M2+A1

3 marks

d. and e.



A2
A2
2 + 2 marks

- f. Reflection in the x –axis, dilation factor of 2 away from the x –axis, dilation factor if $\frac{1}{2}$ away from the y –axis, translation of 2 units to the right and a translation of 1 unit up.

$$\text{equation: } y = -2 \left(\frac{2}{2x-4-3} - 4 \right) + 1 = 9 - \frac{4}{2x-7}$$

M2+A2
4 marks

Question 2

a. $\begin{bmatrix} 0.8 & 0.5 \\ 0.2 & 0.5 \end{bmatrix}^3 \begin{bmatrix} 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 0.722 \\ 0.278 \end{bmatrix}$

$$Pr(\text{candles}) = 0.278$$

M1+A1
2 marks

b. $(0.8)^3 = 0.512$

A1
1 mark

c. $\begin{bmatrix} 0.8 & 0.5 \\ 0.2 & 0.5 \end{bmatrix} \begin{bmatrix} x \\ 1-x \end{bmatrix} = \begin{bmatrix} x \\ 1-x \end{bmatrix}$ or $\frac{0.5}{0.2+0.5}$

$$x = 0.714288$$

$$Pr(\text{coloured light globes}) = 0.7143$$

A1
1 mark

d. i. $\begin{bmatrix} m & -m + 1.3 \\ 1 - m & m - 0.3 \end{bmatrix}^3 \begin{bmatrix} 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 0.671 \\ 0.329 \end{bmatrix}$
 $m = 0.71$

M2+A1
3 marks

ii. $m = 0.71, m - 0.3 = 0.41$

Let C denote buying a candle and L denote buying a coloured light globe

$$Pr(CCC) = 1 \times (0.41)^2 = 0.1681$$

$$Pr(CLC) + Pr(CCL) = 1 \times 0.59 \times 0.29 + 1 \times 0.41 \times 0.59 = 0.413$$

$$Pr(CLL) = 1 \times 0.59 \times 0.71 = 0.4189$$

X	0	1	2
$Pr(X=x)$	0.1681	0.413	0.4189

$$E(X) = 0.413 + 2 \times 0.4189 = 1.251$$

M2+A1
3 marks

e. $Pr(X \geq 2) \geq 0.8$

$$Pr(X = 0) + Pr(X = 1) \leq 0.2$$

$$0.6^n + \binom{n}{1} \times 0.4 \times (0.6)^{n-1} \leq 0.2$$


$$n = 6.4$$

At least 7 coloured light globes needs to be sold.

M2+A1
3 marks

f. $\int_0^{2500} \frac{1}{9.765625 \times 10^{12}} (6.25 \times 10^6 t - t^3) dt = 1$ and $f(t) \geq 0$

M1+A1
2 marks



M1+A1
2 marks

h. $\mu = \int_0^{2500} \frac{t}{9.765625 \times 10^{12}} (6.25 \times 10^6 t - t^3) dt = \frac{4000}{3}$ hours

A1
1 mark

i. $\Pr\left(X > \frac{4000}{3} \mid X \geq 1000\right) = \frac{\Pr(X > \frac{4000}{3})}{\Pr(X \geq 1000)} = \frac{0.51202}{0.7056} = 0.7257$

M1+A1
2 marks

Question 3

a. average value of gradient = $\frac{1}{1.5+1.5} \int_{-1.5}^{1.5} ((x+1)^2(5-4x)) dx = 2.75$

M1+A1
2 marks

b. $f(x) = \int f'(x) dx = -x^4 - x^3 + 3x^2 + 5x + c$
 $f(2) = -16 - 8 + 12 + 10 + c = -2$
 $c = 0$

M1+A1
2 marks

c. $(x+1)^2(5-4x) = 0$
 $x = -1$ point of inflection – gradient is positive before and after the point.
 $x = \frac{5}{4}$ local maximum – gradient changes from positive before the point to negative after the point.

M2+A2
4 marks

d. Area = $-\int_{-1}^0 f(x) dx + \int_0^{1.91964} f(x) dx - \int_{1.91964}^2 f(x) dx = 9.21$. (use the x-int).

M1+A1
2 marks

Question 4

a. $f(x) = |2x^2 - 3x| - 2$


$2x^2 - 3x - 2 = 0$ or $-(2x^2 - 3x) - 2 = 0 \Rightarrow$ no factors.

$(x - 2)(2x + 1) = 0$

$x = 2$ or $x = -\frac{1}{2}$

M1+A2
3 marks

b.



A2
2 marks

Question 5

a. $MS = x$

$\triangle ANC \sim \triangle AMS$

$\frac{AN}{NC} = \frac{AM}{MS}$

$\frac{200}{100} = \frac{AM}{x}$

$AM = 2x$ mm

Area = $2x(200 - 2x) = 400x - 4x^2$ mm²

M2+A2
4 marks

b. $x \in (0, 100)$

$$\frac{dA}{dx} = 400 - 8x = 0$$

$$x = 50 \text{ mm}$$

M1+A1
2 marks

c. $A = 100 \times 100 = 10\,000 \text{ mm}^2$

A1
1 mark