



SECTION 1

| | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|----|----|
| 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 |
| B | A | C | C | A | A | E | E | B | C | A |


| | | | | | | | | | | |
|----|----|----|----|----|----|----|----|----|----|----|
| 12 | 13 | 14 | 15 | 16 | 17 | 18 | 19 | 20 | 21 | 22 |
| A | C | C | B | E | D | B | D | B | E | C |

Q1 $x+3=0, x=-3$ B

Q2 $\log_{\frac{1}{e}}\left(\frac{1}{e^{\sqrt{x}}}\right) = \log_{\frac{1}{e}}\left(\frac{1}{e}\right)^{\sqrt{x}} = \sqrt{x} \log_{\frac{1}{e}}\left(\frac{1}{e}\right) = \sqrt{x}$ A


Q3 $\sin(2x)+\cos(2x)=0, \frac{\sin(2x)}{\cos(2x)}=-1$ where $\cos(2x) \neq 0,$
 $\tan(2x)=-1, 2x=n\pi-\frac{\pi}{4}, x=\frac{(4n-1)\pi}{8}$ C

Q4 $y=-5, x=0; y \rightarrow 0^-, x \rightarrow -\infty; y \rightarrow 2^+, x \rightarrow 7^-;$
 $y \rightarrow \infty, x \rightarrow 2^+$ C




Q6 $f(x)=2(x-a)^2(x-b),$
 $g(x)=2e^x(2x-c-a)^2(2x-c-b)=0$
The two unique solutions are $x=\frac{c+a}{2}$ and $x=\frac{c+b}{2}.$
Since $a > b > 0,$ both solutions are positive if $c > -b.$ A

Q7 $g(x)=f\left(\frac{x}{2}+b\right)=2\left(\left(\frac{x}{2}+b\right)+b\right)^3-a=2\left(\frac{x}{2}+2b\right)^3-a$
 $=2\left(\frac{1}{2}(x+4b)\right)^3-a=\frac{1}{4}(x+4b)^3-a$ E



Q9 The intersection of $y=f(x)$ and $y=f^{-1}(x)$ is on the line $y=x.$
Let $x=e-\log_e(\log_e x), \log_e(\log_e x)=e-x, x=e, \therefore y=e$ B

Q10 $\frac{1}{\sqrt{2}} \leq \sqrt{1-\cos(2x)} \leq 1, \frac{1}{2} \leq 1-\cos(2x) \leq 1,$
 $0 \leq \cos(2x) \leq \frac{1}{2}, \frac{\pi}{3} \leq 2x \leq \frac{\pi}{2}, \frac{\pi}{6} \leq x \leq \frac{\pi}{4}$ C



Q12 Given $g(x)=f^{-1}(x), f(a)=b$ and $f'(a)=\frac{1}{a},$ then
 $g(b)=a$ and $g'(b)=\frac{1}{f'(a)}=a$ A

Q13 Given function $f(x), x \in R,$ then $f(|x|)=\begin{cases} f(x), & x \geq 0 \\ f(-x), & x < 0 \end{cases}.$
 $f(-x)$ for $x < 0$ is the reflection (in the y-axis) of $f(x)$ for $x > 0.$
 $\therefore f'(|x|)=0$ at $(-3,-1), (-1,2), (1,2)$ and $(3,-1)$ C

Q14 $\int_a^b f(x)dx = [F(x)]_a^b = F(b) - F(a) = c$
 $\int_{\frac{a-h}{2}}^{\frac{b-h}{2}} 2f(2x+h)dx = \left[\frac{2F(2x+h)}{2}\right]_{\frac{a-h}{2}}^{\frac{b-h}{2}} = F(b) - F(a) = c$ C

Q15 $\int_0^{\frac{\pi}{8}} g(x)dx = \int_0^{\frac{\pi}{8}} \tan\left(\frac{x}{2} + \frac{\pi}{4}\right)dx = 0.4824$ by CAS
Average value $= \frac{\int_0^{\frac{\pi}{8}} g(x)dx}{\frac{\pi}{8} - 0} \approx \frac{0.4824}{\frac{\pi}{8}} \approx 1.2284$ B

Q16 $f'(x)=\frac{1}{\sqrt{x+10}}, f(x)=\int \frac{1}{\sqrt{x+10}} dx = 2\sqrt{x+10}$
 $bf(a) \approx f(a) + (1.02a - a)f'(a)$
 $bf(a) - f(a) \approx 0.02af'(a), (b-1)f(a) \approx 0.02af'(a)$
 $\therefore (b-1)2\sqrt{a+10} \approx 0.02a \times \frac{1}{\sqrt{a+10}}$
 $b-1 \approx \frac{0.01a}{a+10}, b \approx \frac{1.01a+10}{a+10}$ E

Q17 $\Pr(JandJtogether) = \frac{25!}{6!} = \frac{1}{3}$ D

Q18 $\Pr(\text{1green1blue}) = 2 \times \frac{1}{2} \times \frac{1}{3} = \frac{1}{3}$

Since the two rolls are independent,

$\Pr(\text{green2nd} \mid \text{blue1st}) = \Pr(\text{green2nd}) = \frac{1}{2}$

Q19 Binomial: $np = 12.3$, $np(1-p) = 2.8^2$, $\therefore n \approx 34$

Q20 $B \xrightarrow{\frac{1}{3}} A$, $\therefore B \xrightarrow{\frac{2}{3}} B$

$A \xrightarrow{\frac{1}{4}} A$, $\therefore A \xrightarrow{\frac{3}{4}} B$

$\therefore \Pr(\text{BBAABAB}) = 1 \times \frac{2}{3} \times \frac{1}{3} \times \frac{1}{4} \times \frac{3}{4} \times \frac{1}{3} \times \frac{3}{4} = \frac{1}{96}$

Q21 $\Pr(X > 12.5) = 0.8$, $\Pr(X > 18.5 \mid X > 12.5) = 0.8$

$\therefore \frac{\Pr(X > 18.5 \cap X > 12.5)}{\Pr(X > 12.5)} = 0.8$, $\frac{\Pr(X > 18.5)}{\Pr(X > 12.5)} = 0.8$

$\therefore \Pr(X > 18.5) = 0.8^2 = 0.64$, $\Pr\left(Z > \frac{18.5 - \mu}{\sigma}\right) = 0.64$,


$\therefore \Pr\left(Z < \frac{18.5 - \mu}{\sigma}\right) = 0.36$, $\therefore \frac{18.5 - \mu}{\sigma} = -0.3585$

$\therefore \mu - 0.36\sigma \approx 18.5$

Q22 The graph is a probability density function, $\therefore n \neq \Pr(2)$ C

SECTION 2

Q1a



Q1b $f'(x) = ax^2(x-1)^2$, $\therefore f(x)$ is a degree five polynomial.

| x | < 0 | 0 | $0 < x < 1$ | 1 | > 0 |
|---------|----------|------|-------------|------|----------|
| $f'(x)$ | positive | zero | positive | zero | positive |

The table shows that there is a stationary inflection point at $x = 0$ and another one at $x = 1$.

Q1ci $f'(x) = ax^2(x-1)^2 = a(x^4 - 2x^3 + x^2)$

$\therefore f(x) = a\left(\frac{x^5}{5} - \frac{x^4}{2} + \frac{x^3}{3}\right) + c$, $f(0) = 0$

$\therefore f(x) = a\left(\frac{x^5}{5} - \frac{x^4}{2} + \frac{x^3}{3}\right)$

Q1cii $f(1) = 1$, $\therefore a = 30$

Q1d $f'(x) = 30(x^4 - 2x^3 + x^2)$, $x \in [0, 1]$

$f''(x) = 30(4x^3 - 6x^2 + 2x) = 60(2x^3 - 3x^2 + x)$
 $= 60x(x-1)(2x-1)$

B

Let $f''(x) = 0$ to find the greatest rate of change:

D $60x(x-1)(2x-1) = 0$, $\therefore x = \frac{1}{2}$, $\therefore y = 30\left(\frac{x^5}{5} - \frac{x^4}{2} + \frac{x^3}{3}\right) = \frac{1}{2}$

$\left(\frac{1}{2}, \frac{1}{2}\right)$

B

Q1e Since $\left(\frac{1}{2}, \frac{1}{2}\right)$ is a point having the greatest rate of change, it is an inflection point. \therefore total number of inflection points is 3.

Q1f $g(x) = -2f(-x) + 2$

$f(x)$ undergoes reflection in both axes, a dilation by a factor of 2 parallel to the y-axis and then an upward translation of 2 units.
 $(0, 0) \rightarrow (0, 2)$; $(1, 1) \rightarrow (-1, 0)$

Q1g $g(x) = -2f(-x) + 2$

E

$= -2 \times 30\left(\frac{(-x)^5}{5} - \frac{(-x)^4}{2} + \frac{(-x)^3}{3}\right) + 2$

$= 60\left(\frac{x^5}{5} - \frac{x^4}{2} + \frac{x^3}{3}\right) + 2$, a strictly increasing function (by CAS)

Domain of $f(x)$ is $(-1, 2]$, \therefore domain of $g(x)$ is $[-2, 1)$.

When $x = -2$, $y = -62$; when $x \rightarrow 1$, $y \rightarrow 64$.

\therefore range of $g(x)$ is $[-62, 64)$

Q2a $height = 2\log_e(a+e)$, $width = 2e-a$,

$area A = (2e-a) \times 2\log_e(a+e) = 2(2e-a)\log_e(a+e)$

Q2bi By the product rule:


$\frac{dA}{da} = 2(2e-a) \times \frac{1}{a+e} + (-2)\log_e(a+e) = \frac{2(2e-a)}{a+e} - 2\log_e(a+e)$

Q2bii Maximum cross-sectional area \Rightarrow maximum volume

Let $\frac{dA}{da} = 0$, $\therefore \frac{2(2e-a)}{a+e} - 2\log_e(a+e) = 0$,

$\therefore 2\log_e(a+e) = \frac{2(2e-a)}{a+e}$, $height = \frac{2(2e-a)}{a+e}$

Q2biii



Q2c The tank has maximum volume when $a = 0.8645$

$$\text{Volume of water } V = 8(2e - 0.8645)h,$$

$$\frac{dV}{dh} = 8(2e - 0.8645), \quad \frac{dV}{dt} = 90 - 36h = 54 \quad \text{when } h = 1$$

$$\frac{dV}{dt} = \frac{dV}{dh} \times \frac{dh}{dt}, \quad \therefore 54 = 8(2e - 0.8645) \frac{dh}{dt},$$

$$\frac{dh}{dt} = \frac{54}{8(2e - 0.8645)} \approx 1.48 \text{ m/h}$$

$$\text{Q2d } \frac{dt}{dh} = \frac{1}{2.5 - h}, \quad t = \int \frac{1}{2.5 - h} dh = -\log_e |2.5 - h| + c$$

Given at time $t = 0$, $h = 0$, $\therefore c = \log_e 2.5$

$$\therefore t = -\log_e |2.5 - h| + \log_e 2.5$$

$$\text{When } h = 1.25, \quad t = -\log_e 1.25 + \log_e 2.5 = \log_e \frac{2.5}{1.25} = \log_e 2$$

Q2e Volume of water in the tank is maximum when

$$\frac{dV}{dt} = 90 - 36h = 0, \quad h = 2.5$$

$$V_{\max} = 8(2e - 0.8645) \times 2.5 \approx 91 \text{ m}^3$$

Q2f Cross-sectional area of the tunnel

$$= \int_0^{2e} 2 \log_e(x + e) dx \approx 17.92$$

$$\text{Volume of soil} = (17.92 - 11.67) \times 8 \approx 50 \text{ m}^3$$

Q3a $(0,0)$, $a + c = 0$, $\therefore c = -a$.

$$(10, -5), \quad ae^{10b} + c = -5, \quad \therefore e^{10b} = \frac{-5 - c}{a} = \frac{a - 5}{a}.$$

$$(20, -8), \quad ae^{20b} + c = -8, \quad \therefore e^{20b} = \frac{-8 - c}{a} = \frac{a - 8}{a},$$

$$\therefore (e^{10b})^2 = \frac{a - 8}{a}, \quad \therefore \left(\frac{a - 5}{a}\right)^2 = \frac{a - 8}{a}.$$

$$a^2 - 10a + 25 = a^2 - 8a, \quad 2a = 25, \quad \therefore a = \frac{25}{2}, \quad c = -\frac{25}{2} \text{ and}$$

$$e^{10b} = \frac{a - 5}{a} = \frac{3}{5}, \quad b = \frac{1}{10} \log_e \left(\frac{3}{5}\right)$$

$$\text{Q3aii } y = ae^{bx} + c, \quad \frac{dy}{dx} = abe^{bx} = \frac{25}{2} \times \frac{1}{10} \log_e \left(\frac{3}{5}\right) e^{\frac{x}{10} \log_e \left(\frac{3}{5}\right)}$$

$$= \frac{5}{4} \log_e \left(\frac{3}{5}\right) e^{\log_e \left(\frac{3}{5}\right) \frac{x}{10}} = \left[\frac{5}{4} \log_e \left(\frac{3}{5}\right)\right] \left(\frac{3}{5}\right)^{\frac{x}{10}}$$

Q3aiii When $x = 10$,

$$y = \left[\frac{5}{4} \log_e \left(\frac{3}{5}\right)\right] \left(\frac{3}{5}\right)^{\frac{x}{10}} = \left[\frac{5}{4} \log_e \left(\frac{3}{5}\right)\right] \left(\frac{3}{5}\right) \approx -0.38$$

Q3bi When $t = 6$, $y = 3 \sin\left(\frac{\pi}{6}\right) - 5 = 3 \sin \pi - 5 = -5$. The sea

level is 5 m below the house.

Q3bii When $t = 6$, $y = -5$ and $\therefore x = 10$

\therefore the horizontal distance between the house and the water edge is $2 + 10 = 12$ m

$$\text{Q3c When } t = 6, \quad \frac{dy}{dt} = \frac{\pi}{2} \cos\left(\frac{\pi}{6}\right) = \frac{\pi}{2} \cos \pi = -\frac{\pi}{2}$$

$$\text{Q3d } \frac{dy}{dt} = \frac{dy}{dx} \times \frac{dx}{dt}, \quad -\frac{\pi}{2} = \frac{3}{4} \log_e \left(\frac{3}{5}\right) \times \frac{dx}{dt}$$

$$\therefore \frac{dx}{dt} = -\frac{2\pi}{3 \log_e \left(\frac{3}{5}\right)}, \quad \therefore \text{the receding rate is } \frac{2\pi}{3 \log_e \left(\frac{3}{5}\right)}.$$

Q4a $\Pr(W > 52)$, by CAS *normalcdf*(52, e^{99} , 65, 8) ≈ 0.9479

$$\text{Q4b } \Pr(W > 52 | W > 45) = \frac{\Pr(W > 52)}{\Pr(W > 45)} \approx 0.9538$$

Q4c Average price (\$) per dozen

$$= \frac{\Pr(52 < W < 60)}{\Pr(W > 52)} \times 1.00 + \frac{\Pr(60 < W < 68)}{\Pr(W > 52)} \times 1.10 + \frac{\Pr(W > 68)}{\Pr(W > 52)} \times 1.20 \approx 1.11$$

Q4d Binomial: $n = 12$, success means $65 \leq W < 68$,

$$p = \frac{\Pr(65 < W < 68)}{\Pr(60 < W < 68)} = \frac{0.14617}{0.380184} \approx 0.384472$$

$\Pr(X \geq 6)$ by CAS *binomialcdf*(12, 0.384472, 6, 12) ≈ 0.29

Q4e The average weight of each of the four eggs is less than

$$\frac{250}{4} = 62.5.$$

$$\Pr(\text{total} < 250) = \frac{\Pr(60 < W < 62.5)}{\Pr(60 < W < 68)} = \frac{0.111345}{0.380184} \approx 0.29$$

Q4f Probability density function $f(x) = \frac{1}{8\sqrt{2\pi}} e^{-\frac{1}{2}\left(\frac{x-65}{8}\right)^2}$

$$\text{Mean weight} = 12 \times \frac{\int_{68}^{\infty} xf(x) dx}{\Pr(W > 68)} \approx 12 \times \frac{25.97381}{0.35383} \approx 881$$

Q4g $\Pr(W < 45) = 0.05$ and $\Pr(W > 75) = 0.10$

$$\therefore \Pr\left(Z < \frac{45 - \mu}{\sigma}\right) = 0.05 \quad \text{and} \quad \Pr\left(Z < \frac{75 - \mu}{\sigma}\right) = 0.90$$

$$\therefore \frac{45 - \mu}{\sigma} \approx -1.6449 \quad \text{and} \quad \frac{75 - \mu}{\sigma} \approx 1.2816$$

$$\therefore \mu \approx 61.86 \text{ grams}, \quad \sigma = 10.25 \text{ grams}$$

Q4h $\Pr(ABBBBBA) + \Pr(ABBBBAB) + \Pr(ABBBABB)$

$$+ \Pr(ABBABBB) + \Pr(ABABBBB) + \Pr(AABBBBB)$$

$$= (1)(0.64)(0.45)^4(0.55) + (1)(0.64)(0.45)^3(0.55)(0.64)$$

$$+ (1)(0.64)(0.45)^2(0.55)(0.64)(0.45) + (1)(0.64)(0.45)(0.55)(0.64)(0.45)^2$$

$$+ (1)(0.64)(0.55)(0.64)(0.45)^3 + (1)(0.36)(0.64)(0.45)^4 \approx 0.11$$

Please inform mathline@itute.com re conceptual, mathematical and/or typing errors