

2014 Trial Examination

THIS BOX IS FOR ILLUSTRATIVE PURPOSES ONLY	
]

	STUDEN	L NOMBE	K			_	Letter
Figures							
Words							_

MATHEMATICAL METHODS (CAS)

Units 3 & 4 – Written examination 1

Reading time: 15 minutes Writing time: 1 hour

QUESTION & ANSWER BOOK

Structure of book

	D 11 4 C 1 0 C 0 C 1 1	
Number of questions	Number of questions to be answered	Number of marks
8	8	40
		Total 40

- Students are permitted to bring into the examination room: pens, pencils, highlighters, erasers, sharpeners and rulers.
- Students are NOT permitted to bring into the examination room: notes of any kind, blank sheets of paper, white out liquid/tape or a calculator of any type.

Materials supplied

- Question and answer book of 9 pages.
- Working space is provided throughout the book.

Instructions

- Print your name in the space provided on the top of this page.
- All written responses must be in English.

Students are NOT permitted to bring mobile phones and/or any other unauthorised electronic communication devices into the examination room.

© TSSM 2014 Page 1 of 9

Instructions

Answer **all** questions in the spaces provided.

A decimal approximation will not be accepted if an exact answer is required to a question. In questions where more than one mark is available, appropriate working must be shown. Unless otherwise indicated, the diagrams in this book are **not** drawn to scale.

Question 1(6 marks)

If $f(x) = 2x^2 log_e(3x)$, find $f'(x)$.	2 marks
Find $f'\left(\frac{1}{3}\right)$.	1 mark
Find the value(s) of x for which $f'(x) = 0$.	3 marks
	Find $f'\left(\frac{1}{3}\right)$.

© TSSM 2014 Page 2 of 9

Question 2 (6 marks)

Find the derivative of $(2x - 3)\sin(x)$.	2 mark
Hence, find an anti-derivative of $x\cos(x)$.	4 mark

TURN OVER

© TSSM 2014 Page 3 of 9

Qu	nestion 3 (2 marks)
So	lve the equation $2\cos\left(\frac{x}{3}\right) + 1 = 0$, $x \in [3\pi, 6\pi]$
	
Qu	testion 4 (9 marks)
a.	Find the value of k for which $4e^{2x} - ke^x + 9 = 0$ has a unique solution. 2 marks
	2
b.	Find the unique solution for the equation $4e^{2x} - ke^x + 9 = 0$ for the positive value of k found in part a .

Question 4 - continued

© TSSM 2014 Page 4 of 9

2014 MATHMETH (CAS) EXAM 1

 $Let g(x) = 2e^x - 3$

c.	Find the inverse, $g^{-1}(x)$, of the function $g(x)$.	2 marks	
d.	State the domain and range of $g^{-1}(x)$.	2 marks	

TURN OVER

2014 MATHMETH (CAS) EXAM 1

Question 5 (4 marks) Let $f: R \to R$, $f(x) = 4x + 9$.	
The average value of f on the interval $[0, a]$ is $\frac{1}{a}$,	where a is a real constant.
Find all possible values of <i>a</i> .	

© TSSM 2014 Page 6 of 9

Question 6 (6 marks)

The probability distribution of a discrete random variable, X, is given by the table below.

X	0	1	2	3	4
Pr(X)	1	k	k	3	2
	$\overline{2k}$	$\overline{10}$	$\overline{10}$	$\overline{10k}$	$\overline{5k}$

a. Show that $k=3$	or <i>k</i>	= 2
---------------------------	-------------	-----

3 marks

Let k = 3.

b.	Calculate	E(X)
----	-----------	------

2 marks

c.	Find $Pr(X \leq E(X))$.	1 mark

TURN OVER


© TSSM 2014 Page 7 of 9

2014 MATHMETH (CAS) EXAM 1

Question 7 (4 marks)

Let
$$f:[0, \infty) \to R$$
, $f(x) = 2e^{-\frac{x}{3}}$

Let S be the point on the graph of f on the y-axis and let T be the point on the graph of f with the y-coordinate $\frac{1}{2}$.



Find the area of the region bounded by the graph of f and the line segment ST.

© TSSM 2014 Page 8 of 9

Question 8 (3 marks)

A continuous random variable, X, has a probability density function

$$f(x) = \begin{cases} \frac{1}{2}\sin(x) & 0 \le x \le \pi \\ 0 & elsewhere \end{cases}$$

a.	Find the median of X .	2 marks
b.	Find the mode of X .	1 mark

END OF QUESTION AND ANSWER BOOK

© TSSM 2014 Page 9 of 9