

**SPECIALIST MATHS EXAM 2 SOLUTIONS****Question 1**

a.  $[0, 0.8]$

A1

b.  $f(x) = 1 - (4x^2 + 1)^{-1}$

$$f'(x) = 0 + (4x^2 + 1)^{-2} \times 8x$$

$$= \frac{8x}{(4x^2 + 1)^2}$$

A1

$$f''(x) = \frac{8(4x^2 + 1)^2 - 8x \times 2(4x^2 + 1) \times 8x}{(4x^2 + 1)^4}$$

At point of greatest slope  $f''(x) = 0$ 

$$0 = \frac{8(4x^2 + 1)[(4x^2 + 1) - 16x^2]}{(4x^2 + 1)^4}$$

M1

$$0 = 4x^2 + 1 - 16x^2$$

$$12x^2 = 1$$

$$x = \frac{1}{\sqrt{12}} = \frac{\sqrt{12}}{12} = \frac{\sqrt{3}}{6}$$

(positive root only, since domain is  $[0, 1]$ ) A1

$$y = 1 - \frac{1}{4\left(\frac{\sqrt{3}}{6}\right)^2 + 1} = \frac{1}{4}$$

M1

∴ The point of greatest slope is at  $\left(\frac{\sqrt{3}}{6}, \frac{1}{4}\right)$ .

c.  $y = 1 - \frac{1}{4x^2 + 1}$

Transpose to find  $x^2$ 

$$4x^2 + 1 = \frac{1}{1-y}$$

$$x^2 = \frac{1}{4}\left(\frac{1}{1-y} - 1\right)$$

A1

$$V = \pi \int_0^{0.8} x^2 dy$$

$$V = \frac{\pi}{4} \int_0^{0.8} \left(\frac{1}{1-y} - 1\right) dy$$

M1

$$= \frac{\pi}{4} \left[ -\log_e(1-y) - y \right]_0^{0.8}$$

$$= -\frac{\pi}{4} [\log_e(1-0.8) + 0.8 - \log_e(1-0) - 0]$$

M1

$$= 0.636 \text{ m}^3$$

H1

d.  $\frac{dV}{dt} = \frac{dV}{dy} \times \frac{dy}{dt}$

$$0.012 = \frac{\pi}{4} \left( \frac{1}{1-y} - 1 \right) \times \frac{dy}{dt}$$

M1

$$\left( \frac{1}{1-y} - 1 \right) \frac{dy}{dt} = \frac{0.012 \times 4}{\pi}$$

At point of greatest slope,  $y = \frac{1}{4}$ 

$$\left( \frac{1}{1-\frac{1}{4}} - 1 \right) \frac{dy}{dt} = \frac{0.012 \times 4}{\pi}$$

M1

$$\left( \frac{4}{3} - 1 \right) \times \frac{dy}{dt} = \frac{0.012 \times 4}{\pi}$$


$$\frac{dy}{dt} = \frac{0.012 \times 4}{\pi} \times \frac{3}{1}$$

$$\frac{dy}{dt} = \frac{0.144}{\pi} \text{ m/min}$$

A1

**Question 2**

a.


**A1 Shape**  
**A1 Endpoints**

b. i

$$f(x) = x \cos^{-1}(2x)$$

$$f'(x) = 1 \cdot \cos^{-1}(2x) + x \cdot \frac{-1}{\sqrt{1-(2x)^2}} \times 2$$

M1

$$f'(x) = \cos^{-1}(2x) - \frac{2x}{\sqrt{1-4x^2}}$$

A1

$$\text{ii. } \left\{ x : -\frac{1}{2} < x < \frac{1}{2} \right\} \quad \text{or} \quad \left( -\frac{1}{2}, \frac{1}{2} \right)$$

A1

c.  $g(x) = (1 - 4x^2)^{\frac{1}{2}}$

$$g'(x) = \frac{1}{2}(1 - 4x^2)^{-\frac{1}{2}}(-8x) \quad \mathbf{M1}$$

$$g'(x) = \frac{-4x}{\sqrt{1 - 4x^2}}$$

$$\Rightarrow \left\{ x : -\frac{1}{2} < x < \frac{1}{2} \right\} \quad \mathbf{A1}$$

d. i.  $f'(x) = \cos^{-1}(2x) - \frac{2x}{\sqrt{1 - 4x^2}}$

$$f'(x) = \cos^{-1}(2x) + \frac{1}{2} \times \frac{-4x}{\sqrt{1 - 4x^2}}$$

$$f'(x) = \cos^{-1}(2x) + \frac{1}{2}g'(x) \quad \mathbf{M1}$$

$$\cos^{-1}(2x) = f'(x) - \frac{1}{2}g'(x)$$

$$3\cos^{-1}(2x) = 3\left[f'(x) - \frac{1}{2}g'(x)\right] \quad \mathbf{A1}$$

ii.  $\int_0^{\frac{1}{2}} 3\cos^{-1}(2x) dx$

$$= 3 \int_0^{\frac{1}{2}} f'(x) - \frac{1}{2}g'(x) dx$$

$$= 3 \left[ f(x) - \frac{1}{2}g(x) \right]_0^{\frac{1}{2}} \quad \mathbf{M1}$$

$$= 3 \left[ x\cos^{-1}(2x) - \frac{1}{2}\sqrt{1 - 4x^2} \right]_0^{\frac{1}{2}} \quad \mathbf{M1}$$

$$= 3 \left[ \frac{1}{4}\cos^{-1}(2 \times \frac{1}{4}) - \frac{1}{2}\sqrt{1 - 4\left(\frac{1}{4}\right)^2} - 0 + \frac{1}{2} \right]$$

$$= 3 \left[ \frac{1}{4}\cos^{-1}\left(\frac{1}{2}\right) - \frac{1}{2}\sqrt{\frac{3}{4} + \frac{1}{2}} \right]$$

$$= 3 \left[ \frac{1}{4} \times \frac{\pi}{3} - \frac{\sqrt{3}}{4} + \frac{1}{2} \right]$$

$$= 3 \left[ \frac{\pi}{12} - \frac{3\sqrt{3}}{12} + \frac{6}{12} \right]$$

$$= \frac{\pi - 3\sqrt{3} + 6}{4} \quad \mathbf{A1}$$

**Question 3**

a.




fig 1




fig 2

fig 1	fig 2
$r = \sqrt{(\sqrt{3})^2 + 1^2} = 2$	$r = \sqrt{(-\sqrt{3})^2 + 1^2} = 2$
$\sin \theta = \frac{1}{2}$ or $\tan \theta = \frac{1}{\sqrt{3}}$	$\cos \phi = \frac{\pi}{6}$ or $\tan \phi = \frac{\sqrt{3}}{1}$
$\theta = \frac{\pi}{6}$	$\phi = \frac{\pi}{3}$
$u = 2\text{cis}\frac{\pi}{6}$	$v = 2\text{cis}\left(-\frac{\pi}{3}\right)$


b.  $uv = (\sqrt{3} + i)(1 - \sqrt{3}i)$

$$= \sqrt{3} - 3i + i - \sqrt{3}i^2$$

$$= 2\sqrt{3} - 2i$$

A1


c.



$i^2 u = -1(\sqrt{3} + i)$	$r = \sqrt{(-\sqrt{3})^2 + (-1)^2} = 2$
$i^2 u = -\sqrt{3} - i$	$\tan \theta = \frac{1}{\sqrt{3}}$
$i^2 u = 2\text{cis}\left(-\frac{5\pi}{6}\right)$	$\theta = \frac{\pi}{6}$

A1

d.



e.  $i^2 u = 2\text{cis}\left(-\frac{5\pi}{6}\right)$  from c.  
 $i^3 v = i^2 i \times (1 - \sqrt{3}i)$   
 $= -i(1 - \sqrt{3}i)$   
 $= -\sqrt{3} - i$   
 $= 2\text{cis}\left(-\frac{5\pi}{6}\right)$  A1

Multiplication by  $i$  represents an anticlockwise rotation of  $90^\circ$  or  $\frac{\pi}{2}$ . Since  $u$  and  $v$  are perpendicular, if  $u$  is rotated by  $180^\circ$  ( $i^2$ ) and  $v$  rotated by  $270^\circ$  ( $i^3$ ) then the two complex numbers will coincide and hence will be equal. A1

**Question 4**

a.  $\frac{dN}{dt} = kN$

$$t = \frac{1}{k} \int \frac{1}{N} dN + c$$

$$t = \frac{1}{k} \log_e N + c$$

When  $t = 0, N = 700$

$$c = -\frac{1}{k} \log_e 700$$

$$t = \frac{1}{k} (\log_e N - \log_e 700)$$

$$t = \frac{1}{k} \log_e \left( \frac{N}{700} \right)$$

$$e^{kt} = \frac{N}{700}$$

$$N = 700e^{kt}$$

M1

A1

b. When  $t = 2, N = 550$

$$550 = 700e^{2k}$$

$$k = \frac{1}{2} \log_e \left( \frac{550}{700} \right)$$

$$k = -0.12$$

A1

c.  $700e^{-0.12t} < 50$

$$t > -\frac{1}{0.12} \log_e \left( \frac{50}{700} \right) = 22 \text{ years}$$

A1

d.  $\frac{dN}{dt} = P + mN$

$$t = \frac{1}{m} \int \frac{m}{P + mN} dN$$

$$t = \frac{1}{m} \log_e (P + mN) + c$$

When  $t = 3, N = 488$

$$c = 3 - \frac{1}{m} \log_e (P + 488m)$$

$$t = \frac{1}{m} \log_e (P + mN) + 3 - \frac{1}{m} \log_e (P + 488m)$$

$$t - 3 = \frac{1}{m} \log_e \left( \frac{P + mN}{P + 488m} \right)$$

$$e^{m(t-3)} = \frac{P + mN}{P + 488m}$$

$$N = \frac{(P + 488m)e^{m(t-3)} - P}{m}$$

M1

A1

e. i. If  $P = 60$  and  $m = -0.05$

$$N = \frac{(P + 488m)e^{m(t-3)} - P}{m}$$

$$N = \frac{35.6e^{-0.05(t-3)} - 60}{-0.05}$$

When  $t = 8$   $N = 645$  penguins A1

ii. As  $t \rightarrow \infty$ ,  $N \rightarrow \frac{-60}{-0.05} = 1200$  penguins

A1

iii.  $\frac{(P + 488m)e^{m(t-3)} - P}{m} < 800$


As  $t \rightarrow \infty$   $e^{m(t-3)} \rightarrow 0$  since  $m < 0$  A1

$$\Rightarrow \frac{-P}{m} < 800$$

$$-P > 800m \quad \text{since } m < 0$$

$$\therefore P < -800m \quad \text{A1}$$

### Question 5



a.  $N = mg \cos \theta$

$$= 35 \times 9.8 \cos 60^\circ$$

$$= 171.5 \text{ newtons}$$

A1

b. i.  $u = 0 \text{ m/s}, s = 5 \text{ metres}, v = 8 \text{ m/s}$

$$v^2 - u^2 = 2as$$

$$a = \frac{v^2 - u^2}{2s}$$

$$= \frac{64}{10}$$

$$= 6.4 \text{ m/s}^2$$

A1

b. ii. From  $R = ma$

$$mg \sin \theta - F = ma, \text{ where } F = \mu N$$

$$mg \sin \theta - \mu mg \cos \theta = ma$$

M1

$$g \sin \theta - \mu g \cos \theta = a$$

$$\mu = \frac{g \sin \theta - a}{g \cos \theta}$$

$$= \frac{9.8 \sin 60 - 6.4}{9.8 \cos 60}$$

$$= 0.43$$

A1

c. i.  $p = mv$

$$= 35 \times 8$$

$$= 280 \text{ kg m/s}$$

A1

ii.  $(35 + 3)v = 280$

$$v = \frac{140}{19} \text{ m/s}$$

A1

d.  $u = \frac{140}{19} \text{ m/s}, t = 1.4 \text{ seconds}, s = 16 \text{ metres}$

$$s = ut + \frac{1}{2}at^2 \quad \text{M1}$$

$$16 = \frac{140}{19} \times 1.4 + \frac{1}{2}a \times 1.4^2$$

$$a = 5.8 \text{ m/s}^2$$

$$\mu = \frac{g \sin \theta - a}{g \cos \theta}$$

$$= \frac{9.8 \sin 60 - 5.8}{9.8 \cos 60}$$

$$= 0.55$$

M1

e.  $u = \frac{140}{19} \text{ m/s}, t = 1.4 \text{ seconds}, s = 16 \text{ metres}$

$$s = \frac{(u + v)t}{2}$$


$$v = \frac{2s}{t} - u$$

$$v = \frac{2 \times 16}{1.4} - \frac{140}{19}$$

$$= 15.5 \text{ m/s}$$

A1

f.



$$\sin 60 = \frac{y}{15.49} \quad y = 13.41 \quad \mathbf{M1}$$

$$\cos 60 = \frac{x}{15.49} \quad x = 7.74$$

$$\underset{\sim}{v} = \underset{\sim}{7.74} \underset{\sim}{i} - \underset{\sim}{13.41} \underset{\sim}{j} \quad \mathbf{A1}$$

$$\mathbf{g.} \quad \underset{\sim}{v} = -gt \underset{\sim}{j} + \underset{\sim}{c} \quad \mathbf{M1}$$

$$\underset{\sim}{t} = 0, \underset{\sim}{v} = \underset{\sim}{7.74} \underset{\sim}{i} - \underset{\sim}{13.41} \underset{\sim}{j}$$

$$\underset{\sim}{c} = \underset{\sim}{7.74} \underset{\sim}{i} - \underset{\sim}{13.41} \underset{\sim}{j}$$

$$\underset{\sim}{v} = \underset{\sim}{7.74} \underset{\sim}{i} - (\underset{\sim}{13.41} + gt) \underset{\sim}{j} \quad \mathbf{A1}$$

$$\mathbf{h. i} \quad \text{speed} = \left| \underset{\sim}{v} \right| = \sqrt{7.74^2 + (13.41 + 9.8 \times 1.5)^2} \\ = 29.2 \text{ m/s} \quad \mathbf{A1}$$

- ii The horizontal component of the velocity is constant at 7.74 m/s.

$$\begin{aligned} v &= \frac{d}{t} \\ d &= vt \\ &= 7.74 \times 1.5 \\ &= 11.6 \text{ metres} \quad \mathbf{A1} \end{aligned}$$