

Q1  $\int \frac{6+x}{x^2+4} dx = \int \left( \frac{6}{x^2+4} + \frac{x}{x^2+4} \right) dx$   
 $\int \frac{6}{x^2+4} dx + \int \frac{x}{x^2+4} dx = 3 \int \frac{2}{x^2+4} dx + \frac{1}{2} \int \frac{1}{u} du$   $u = x^2 + 4$   
 $= 3 \tan^{-1} \left( \frac{x}{2} \right) + \frac{1}{2} \log_e (x^2 + 4) + C$

Q2  $2 \cos x = \sqrt{3} \cot x$ ,  $2 \cos x - \sqrt{3} \cot x = 0$   
 $2 \cos x - \sqrt{3} \frac{\cos x}{\sin x} = 0$ ,  $\left( 2 - \frac{\sqrt{3}}{\sin x} \right) \cos x = 0$   
 $\therefore 2 - \frac{\sqrt{3}}{\sin x} = 0$ , i.e.  $\sin x = \frac{\sqrt{3}}{2}$ ,  $x = \left( 2n + \frac{1}{2} \mp \frac{1}{6} \right) \pi$

OR  $\cos x = 0$ , i.e.  $x = \left( n + \frac{1}{2} \right) \pi$ , where  $n = 0, \pm 1, \pm 2, \dots$

Q3a Given  $z = 2cis\left(\frac{2\pi}{3}\right) = -1 + i\sqrt{3}$  is a root of the equation

$z^3 - z^2 - 2z - 12 = 0$  which has real coefficients,


$\therefore z = -1 - i\sqrt{3}$  is also a root.

Since  $(z - \alpha)(z - \beta)(z - \gamma) = z^3 - z^2 - 2z - 12$  where


$\alpha = -1 + i\sqrt{3}$  and  $\beta = -1 - i\sqrt{3}$  and  $\gamma$  is the third root,

$\therefore \alpha\beta\gamma = 12$ ,  $(-1 + i\sqrt{3})(-1 - i\sqrt{3})\gamma = 12$ ,  $\therefore 4\gamma = 12$ ,  $\gamma = 3$ .

Q3b



Q4a



Q4b Without the crate leaving the floor, maximum tension  $T_{\max}$  occurs when the normal reaction force  $N \rightarrow 0$ ,  
 $T_{\max} \sin 30^\circ - 50g = 0$ ,  $\therefore T_{\max} = 100g = 980 \text{ N}$

Q4c On the point of moving:

$T \sin 30^\circ + N - 50g = 0$  ..... (1)

and  $T \cos 30^\circ - \mu N = 0$  ..... (2)

From (2),  $N = \frac{T \cos 30^\circ}{\mu} = \frac{5\sqrt{3}T}{2}$  ..... (3)

Substitute (3) in (1):  $\frac{T}{2} + \frac{5\sqrt{3}T}{2} - 50g = 0$ ,  $\therefore (1 + 5\sqrt{3})T = 100g$

$T = \frac{100g}{1 + 5\sqrt{3}} \text{ N}$

Q5  $y = \tan^{-1}(2x)$ ,  $\frac{dy}{dx} = \frac{2}{1+(2x)^2} = \frac{2}{1+4x^2} = 2(1+4x^2)^{-1}$

$\frac{d^2y}{dx^2} = -2(1+4x^2)^{-2}(8x) = -\frac{16x}{(1+4x^2)^2} = -4x \left( \frac{2}{1+4x^2} \right)^2$

Comparing with  $\frac{d^2y}{dx^2} = ax \left( \frac{2}{1+4x^2} \right)^2$ ,  $a = -4$

Q6  $xy^2 + y + (\log_e(x-2))^2 = 14$

Implicit differentiation:  $\frac{d}{dx}(xy^2) + \frac{dy}{dx} + \frac{d}{dx}(\log_e(x-2))^2 = 0$

$\therefore y^2 + 2xy \frac{dy}{dx} + \frac{dy}{dx} + \frac{2 \log_e(x-2)}{x-2} = 0$

At (3,2),  $4 + 13 \frac{dy}{dx} = 0$ ,  $\therefore \frac{dy}{dx} = -\frac{4}{13}$

Q7  $y = (x-1)\sqrt{2-x}$ ,  $1 \leq x \leq 2$

Let  $y = 0$  to find the x-intercepts:  $(x-1)\sqrt{2-x} = 0$ ,  $x = 1, 2$   
 $y > 0$  for  $1 \leq x \leq 2$

Area of the region enclosed by the curve and the x-axis

$= \int_1^2 (x-1)\sqrt{2-x} dx$

$= \int_1^0 -(1-u)u^{\frac{1}{2}} du$

$= \int_0^1 (1-u)u^{\frac{1}{2}} du$

$= \int_0^1 \left( u^{\frac{1}{2}} - u^{\frac{3}{2}} \right) du$

$= \left[ \frac{2u^{\frac{3}{2}}}{3} - \frac{2u^{\frac{5}{2}}}{5} \right]_0^1 = \frac{2}{3} - \frac{2}{5} = \frac{4}{15}$

Let  $u = 2 - x$ ,  
 $\therefore x - 1 = 1 - u$  and  
 $\frac{du}{dx} = -1$   
 When  $x = 1$ ,  $u = 1$ .  
 When  $x = 2$ ,  $u = 0$ .



Q8  $v = \frac{2x}{\sqrt{1+x^2}}, \frac{1}{2}v^2 = \frac{2x^2}{1+x^2}$

$$a = \frac{d}{dx} \left( \frac{1}{2}v^2 \right) = \frac{d}{dx} \left( \frac{2x^2}{1+x^2} \right) = \frac{(1+x^2)(4x) - (2x^2)(2x)}{(1+x^2)^2}$$

$$= \frac{4x}{(1+x^2)^2}$$

Q9a  $\tilde{r}(t) = (2\sqrt{t^2+2} - t^2)\tilde{i} + (2\sqrt{t^2+2} + 2t)\tilde{j}, t \geq 0$

$$\tilde{v}(t) = \frac{d\tilde{r}}{dt} = \left( \frac{2t}{\sqrt{t^2+2}} - 2t \right)\tilde{i} + \left( \frac{2t}{\sqrt{t^2+2}} + 2 \right)\tilde{j}$$

Q9b At  $t=1, \tilde{v} = \left( \frac{2}{\sqrt{3}} - 2 \right)\tilde{i} + \left( \frac{2}{\sqrt{3}} + 2 \right)\tilde{j}$ ,

and the speed  $|\tilde{v}| = \sqrt{\left( \frac{2}{\sqrt{3}} - 2 \right)^2 + \left( \frac{2}{\sqrt{3}} + 2 \right)^2} = \frac{4\sqrt{6}}{3}$

Q9c  $\frac{dy}{dt} = \frac{dy}{dx} \times \frac{dx}{dt}, \therefore \frac{dy}{dx} = \frac{\frac{dy}{dt}}{\frac{dx}{dt}} = \frac{\frac{2}{\sqrt{3}} + 2}{\frac{2}{\sqrt{3}} - 2} = \frac{1 + \sqrt{3}}{1 - \sqrt{3}}$  at  $t=1$

Q9d At time  $t=0, \tilde{r} = 2\sqrt{2}\tilde{i} + 2\sqrt{2}\tilde{j}$  and makes an angle of  $\frac{\pi}{4}$  with the positive  $x$ -axis, whilst vector  $-\sqrt{3}\tilde{i} + \tilde{j}$  makes an angle of  $\frac{5\pi}{6}$  with the positive  $x$ -axis.

$$\therefore \text{angle between the two vectors} = \frac{5\pi}{6} - \frac{\pi}{4} = \frac{7\pi}{12}$$

Q10ai  $-1 \leq \frac{x}{2} \leq 1, \therefore -2 \leq x \leq 2$ . The maximal domain of

$$f_1(x) = \sin^{-1}\left(\frac{x}{2}\right) \text{ is } [-2, 2].$$

Q10aaii  $25x^2 - 1 > 0, \therefore x < -\frac{1}{5}$  or  $x > \frac{1}{5}$ . The maximal domain

$$\text{of } f_2(x) = \frac{3}{\sqrt{25x^2 - 1}} \text{ is } \left(-\infty, -\frac{1}{5}\right) \cup \left(\frac{1}{5}, \infty\right).$$

Q10aiii  $f(x) = \sin^{-1}\left(\frac{x}{2}\right) + \frac{3}{\sqrt{25x^2 - 1}}$  is defined over the

intersection of the maximal domains of  $f_1(x)$  and  $f_2(x)$ , i.e.

$$\left[-2, -\frac{1}{5}\right) \cup \left(\frac{1}{5}, 2\right].$$

Q10b  $h(x) = \sin^{-1}\left(\frac{x}{2}\right) + \sin^{-1}(3x)$

Let  $\theta = h\left(\frac{1}{4}\right) = \sin^{-1}\left(\frac{1}{8}\right) + \sin^{-1}\left(\frac{3}{4}\right) = \alpha + \beta$  where

$$\alpha = \sin^{-1}\left(\frac{1}{8}\right) \text{ and } \beta = \sin^{-1}\left(\frac{3}{4}\right).$$

$$\therefore \sin \alpha = \frac{1}{8} \text{ and } \sin \beta = \frac{3}{4}$$

$$\therefore \cos \alpha = \frac{3\sqrt{7}}{8} \text{ and } \cos \beta = \frac{\sqrt{7}}{4} \text{ by the identity}$$

$$\sin^2 A + \cos^2 A = 1.$$

$$\therefore \sin \theta = \sin(\alpha + \beta) = \sin \alpha \cos \beta + \cos \alpha \sin \beta$$

$$= \frac{1}{8} \times \frac{\sqrt{7}}{4} + \frac{3\sqrt{7}}{8} \times \frac{3}{4} = \frac{5\sqrt{7}}{16}$$

Please inform [mathline@itute.com](mailto:mathline@itute.com) re conceptual, mathematical and/or typing errors