


D

Free Exam
for 2006-15 VCE study design



Units 3 and 4 Specialist Maths: Exam 1

Practice Exam Question and Answer Booklet

Duration: 15 minutes reading time, 1 hour writing time

Structure of book:

Number of questions	Number of questions to be answered	Number of marks
10	10	40

- Students are permitted to bring into the examination room: pens, pencils, highlighters, erasers and rulers.
- Students are not permitted to bring into the examination room: blank sheets of paper and/or white out liquid/tape.
- No calculator is allowed in this examination.

Materials supplied:

- This question and answer booklet of 10 pages.

Instructions:

- You must complete all questions of the examination.
- Write all your answers in the spaces provided in this booklet.

Instructions

Answer all questions in the spaces provided.

In all questions where a numerical answer is required an exact value must be given unless otherwise specified.


In questions where more than one mark is available, appropriate working must be shown.

Unless otherwise indicated, the diagrams in this book are not drawn to scale.

Questions

Question 1

On the axes provided below, draw a graph of the function $f(x) = \frac{-1}{x^2+2x+9}$, labelling all axes intercepts, asymptotes, and stationary points.



4 marks

Question 2

A quadrilateral has four vertices at points O, A, B and C, with coordinates given by (0, 0), (2, 5), (7, 6) and (5, 1). Using vectors, show that $OABC$ is a parallelogram.

3 marks

Question 3

Let $x^2y + \log_e y + x = 2$. Find $\frac{dy}{dx}$.

2 marks

Question 4

Solve $P(z) = z^2 + 5z + 8 = 0$ for $z \in \mathbb{C}$.

3 marks

Question 5

Find the area enclosed by the graph $\frac{7x+1}{x^2+2x-8}$ and the lines $x = -2$ and $x = 0$.

4 marks

Question 6

Let $x = \sec^2(t)$ and $y = \tan(t)$ for $t \in \{t: 0 \leq t \leq \frac{\pi}{4}\}$

- a. Give y as a function of x .

2 marks

- b. State the domain and range of this function.

2 marks

- c. Graph y in terms of x in the space provided below, giving the coordinates and t values of any intercepts and endpoints.

3 marks
Total: 7 marks

Question 7

Show that $\cosec\left(\frac{5\pi}{12}\right) = \sqrt{6} - \sqrt{2}$.

3 marks

Question 8

Evaluate $\int \frac{\tan^{-1}\left(\frac{x}{5}\right)}{25+x^2} dx$.

3 marks

Question 9

Find the exact value of $\int_0^\pi \cos^2(x)\sin^2(x)dx$.

4 marks

Question 10

A particle's displacement is given by the vector function $\mathbf{r}(t) = t^3 \mathbf{i} + \log_e t \mathbf{j}$, where $t > 0$.

- a. Find a Cartesian equation for the particle's displacement.

1 mark

- b. Find an equation for the speed of the particle at time t .

2 marks

- c. At what time(s) is this speed a minimum?

4 marks

Total: 7 marks

End of Booklet

Looking for solutions? Visit www.engageeducation.org.au/practice-exams

To enrol in one of our Specialist Mathematics lectures head to: www.engageeducation.org.au/lectures

Formula sheet

Mensuration

area of a trapezium $\frac{1}{2}(a + b)h$

curved surface area of a cylinder $2\pi rh$

volume of a cylinder $\pi r^2 h$

volume of a cone $\frac{1}{3}\pi r^2 h$

volume of a pyramid $\frac{1}{3}Ah$

volume of a sphere $\frac{4}{3}\pi r^3$

area of a triangle $\frac{1}{2}bc \sin A$

sine rule $\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$

cosine rule $c^2 = a^2 + b^2 - 2ab \cos C$

Coordinate geometry

ellipse	$\frac{(x-h)^2}{a^2} + \frac{(y-k)^2}{b^2} = 1$	hyperbola	$\frac{(x-h)^2}{a^2} - \frac{(y-k)^2}{b^2} = 1$
---------	---	-----------	---

Circular (trigonometric) functions

$$\cos^2(x) + \sin^2(x) = 1$$

$$1 + \tan^2(x) = \sec^2(x)$$

$$\cot^2(x) + 1 = \operatorname{cosec}^2(x)$$

$$\sin(x+y) = \sin(x)\cos(y) + \cos(x)\sin(y)$$

$$\sin(x-y) = \sin(x)\cos(y) - \cos(x)\sin(y)$$

$$\cos(x+y) = \cos(x)\cos(y) - \sin(x)\sin(y)$$

$$\cos(x-y) = \cos(x)\cos(y) + \sin(x)\sin(y)$$

$$\tan(x+y) = \frac{\tan(x) + \tan(y)}{1 - \tan(x)\tan(y)}$$

$$\tan(x-y) = \frac{\tan(x) - \tan(y)}{1 + \tan(x)\tan(y)}$$

$$\cos(2x) = \cos^2(x) - \sin^2(x) = 2\cos^2(x) - 1 = 1 - 2\sin^2(x)$$

$$\sin(2x) = 2\sin(x)\cos(x)$$

$$\tan(2x) = \frac{2\tan(x)}{1 - \tan^2(x)}$$

function	\sin^{-1}	\cos^{-1}	\tan^{-1}
domain	$[-1, 1]$	$[-1, 1]$	\mathbb{R}
range	$[-\frac{\pi}{2}, \frac{\pi}{2}]$	$[0, \pi]$	$(-\frac{\pi}{2}, \frac{\pi}{2})$

Algebra (complex numbers)

$$z = x + yi = r(\cos \theta + i \sin \theta) = r \operatorname{cis} \theta$$

$$z^n = r^n \operatorname{cis}(n\theta) \text{ (de Moivre's theorem)}$$

$$|z| = \sqrt{x^2 + y^2} = r$$

$$-\pi < \operatorname{Arg} z \leq \pi$$

$$z_1 z_2 = r_1 r_2 \operatorname{cis}(\theta_1 + \theta_2)$$

$$\frac{z_1}{z_2} = \frac{r_1}{r_2} \operatorname{cis}(\theta_1 - \theta_2)$$

Calculus

$$\frac{d}{dx}(x^n) = nx^{n-1}$$

$$\int x^n dx = \frac{1}{n+1} x^{n+1} + c, n \neq -1$$

$$\frac{d}{dx}(e^{ax}) = ae^{ax}$$

$$\int e^{ax} dx = \frac{1}{a} e^{ax} + c$$

$$\frac{d}{dx}(\log_e x) = \frac{1}{x}$$

$$\int \frac{1}{x} dx = \log_e |x| + c$$

$$\frac{d}{dx}(\sin(ax)) = a \cos(ax)$$

$$\int \sin(ax) dx = -\frac{1}{a} \cos(ax) + c$$

$$\frac{d}{dx}(\cos(ax)) = -a \sin(ax)$$

$$\int \cos(ax) dx = \frac{1}{a} \sin(ax) + c$$

$$\frac{d}{dx}(\tan(ax)) = \frac{a}{\cos^2(ax)} = a \sec^2(ax)$$

$$\int \sec^2(ax) dx = \frac{1}{a} \tan(ax) + c$$

$$\frac{d}{dx}(\sin^{-1}(x)) = \frac{1}{\sqrt{1-x^2}}$$

$$\int \frac{1}{\sqrt{a^2-x^2}} dx = \sin^{-1}\left(\frac{x}{a}\right) + c, a > 0$$

$$\frac{d}{dx}(\cos^{-1}(x)) = -\frac{1}{\sqrt{1-x^2}}$$

$$\int \frac{-1}{\sqrt{a^2-x^2}} dx = \cos^{-1}\left(\frac{x}{a}\right) + c, a > 0$$

$$\frac{d}{dx}(\tan^{-1}(x)) = \frac{1}{1+x^2}$$

$$\int \frac{a}{a^2+x^2} dx = \tan^{-1}\left(\frac{x}{a}\right) + c$$

product rule

$$\frac{d}{dx}(uv) = u \frac{dv}{dx} + v \frac{du}{dx}$$

quotient rule

$$\frac{d}{dx}\left(\frac{u}{v}\right) = \frac{\left(v \frac{du}{dx} - u \frac{dv}{dx}\right)}{v^2}$$

chain rule

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx}$$

Euler's method

$$\text{If } \frac{dy}{dx} = f(x), x_0 = a \text{ and } y_0 = a, \text{ then } y_{n+1} = y_n + hf(x_n)$$

acceleration

$$a = \frac{d^2x}{dt^2} = \frac{dv}{dt} = v \frac{dv}{dx} = \frac{d}{dx}\left(\frac{1}{2}v^2\right)$$

constant (uniform) acceleration

$$v = u + at, s = ut + \frac{1}{2}at^2, v^2 = u^2 + 2as, s = \frac{1}{2}(u+v)t$$

Vectors in two and three dimensions

$$\mathbf{r} = x\mathbf{i} + y\mathbf{j} + z\mathbf{k}$$

$$|\mathbf{r}| = \sqrt{x^2 + y^2 + z^2} = r$$

$$\mathbf{r} = \frac{d\mathbf{r}}{dt} = \frac{dx}{dt}\mathbf{i} + \frac{dy}{dt}\mathbf{j} + \frac{dz}{dt}\mathbf{k}$$

$$\mathbf{r}_1 \cdot \mathbf{r}_2 = r_1 r_2 \cos \theta = x_1 x_2 + y_1 y_2 + z_1 z_2$$

Mechanics

momentum

$$\mathbf{p} = m\mathbf{v}$$

equation of motion

$$\mathbf{R} = m\mathbf{a}$$

friction

$$F \leq \mu N$$