



SPECIALIST MATHEMATICS Units 3 & 4

Written examination 2 Solutions

SECTION 1

Question	Answer	Solution
1	C	$f(x) = \frac{x}{x^2 - 4x + 3}$ $= \frac{x}{(x-1)(x-3)}$
2	D	<p>Let $u = \sqrt{x-1}$</p> $\Rightarrow x-1 = u^2$ $x = u^2 + 1$ $dx = 2udu$ $\int_2^3 x\sqrt{x-1} dx = \int_1^{\sqrt{2}} (u^2 + 1) \times u \cdot 2udu$ $= \int_1^{\sqrt{2}} 2u^2(u^2 + 1) du$ <p>The terminals of the integral become: When $x = 2$, $u = 1$. When $x = 3$, $u = \sqrt{2}$.</p>
3	B	$V = \rho \int_0^1 (x\sqrt{x})^2 dx$ $= \frac{\rho}{4}$
4	C	<p>The three forces are in equilibrium, $\vec{F}_1 + \vec{F}_2 + \vec{F}_3 = 0$.</p> <p>Use the converse of Pythagoras theorem to show that the triangle formed by the three forces is a right-angled triangle.</p> $ \vec{F}_1 ^2 + \vec{F}_2 ^2 = \vec{F}_3 ^2$ $5^2 + 12^2 = 13^2$ $25 + 144 = 169$

		<p>$169 = 169 \triangleright$ The angle between \vec{F}_1 and \vec{F}_2 is 90°.</p> <p>Use the cosine rule to calculate angle a°.</p> $\cos(a^\circ) = \frac{\left \vec{F}_1\right ^2 + \left \vec{F}_3\right ^2 - \left \vec{F}_2\right ^2}{2\left \vec{F}_1\right \left \vec{F}_3\right }$ $= \frac{25 + 169 - 144}{2 \cdot 5 \cdot 13}$ $= \frac{5}{13}$ $a^\circ = \cos^{-1}\left(\frac{5}{13}\right)$ $= 67.4^\circ$
5	E	$\frac{dy}{dx} = \frac{x}{y}$ $xdx = ydy$ $\int x dx = \int y dy$ $\frac{x^2}{2} + c_1 = \frac{y^2}{2} + c_2$ $x^2 = y^2 + c$, where $c = 2(c_2 - c_1)$ <p>When $x = 0, y = 0 \Rightarrow c = 0$ and $y^2 = x^2$.</p> <p>\Rightarrow first statement is true</p> <p>There are no restrictions on the domain, therefore x can take negative values also.</p> <p>\Rightarrow second statement is false</p> <p>There are no asymptotes for $y = x$.</p> <p>\Rightarrow third statement is false</p> <p>To have a stationary point at $x = 0$, $\frac{dy}{dx} = \frac{x}{y} = 0$. However, when $x = 0, y = 0$, therefore the derivative function is undefined at $x = 0$.</p> <p>\Rightarrow fourth statement is false</p> $\frac{d^2y}{dx^2} = \frac{y - y'x}{y^2}$ $= \frac{y - \frac{x^2}{y}}{y^2}$ $= \frac{y^2 - x^2}{y^3}$ $= 0 \text{ " } x \neq 0 \text{ since } y^2 = x^2$ <p>The second derivative is undefined at $x = 0$.</p> <p>\Rightarrow the fifth statement is false</p>

6	A	<p>Two vectors are linearly dependent if they have the same direction.</p> $\mathbf{a} = k\mathbf{b}, k \in R$ $m\mathbf{i} + n\mathbf{j} = k p\mathbf{i} + k q\mathbf{j}$ $m = kp \text{ and } n = kq$ $\frac{m}{p} = \frac{n}{q} = k$
7	B	<p>$E(3Y - 2X) = 25$ becomes $3E(Y) - 2E(X) = 25$</p> <p>$E(4X - Y) = 18$ becomes $4E(X) - E(Y) = 18$</p> <p>Solve the system of two simultaneous equations for $E(X)$ and $E(Y)$.</p> $E(X) = 7.9, E(Y) = 13.6$ <p>Substitute the values for $E(X)$ and $E(Y)$ into $E(-5X + Y)$.</p> $E(-5X + Y) = -5E(X) + E(Y)$ $= -5 \cdot 7.9 + 13.6$ $= -25.9$
8	D	<p>The maximal domain of $f(x)$ is determined by the intersection between the domains of the two terms of the function.</p> $\frac{x}{1-x} \geq 0 \text{ when } x \in [0, 1)$ $4x \in \left(-\frac{\rho}{2}, \frac{\rho}{2}\right)$ $x \in \left(-\frac{\rho}{8}, \frac{\rho}{8}\right)$ <p>The intersection of the two intervals is $\left[0, \frac{\rho}{8}\right]$.</p>
9	C	<p>$\cos(x) + \sin(x) = a, x \in [0, 2\pi]$</p> <p>Multiply the equation by $\frac{\sqrt{2}}{2}$.</p> $\frac{1}{\sqrt{2}}\cos(x) + \frac{1}{\sqrt{2}}\sin(x) = \frac{1}{\sqrt{2}}a$ $\sin\left(x + \frac{\rho}{4}\right) = \frac{a}{\sqrt{2}} \in [-1, 1]$ <p>$-1 \leq \frac{a}{\sqrt{2}} \leq 1 \dots$ multiply the equation by $\sqrt{2}$</p> $-\sqrt{2} \leq a \leq \sqrt{2}$ $a \in [-\sqrt{2}, \sqrt{2}]$

10	E	$\bar{x} = \frac{103.5 + 108.3}{2} = 105.9$ <p>The z score for the 95% confidence interval is $z = 1.96$.</p> $\bar{x} + z \cdot \frac{s}{\sqrt{n}} = 105.9 + 1.96 \cdot \frac{s}{\sqrt{80}}$ $\Rightarrow 105.9 + 1.96 \cdot \frac{s}{\sqrt{80}} = 108.3$ $s = 10.95217$ $s = 11.0 \text{ mm}$
11	C	<p>Let $u = \frac{1}{5}x^2 + \frac{4}{5}x$</p> $\frac{du}{dx} = \frac{2}{5}x + \frac{4}{5} \Rightarrow du = \left(\frac{2}{5}x + \frac{4}{5}\right)dx$ $du = \frac{2}{5}(x+2)dx$ $\frac{5}{2}du = (x+2)dx$ <p>When $x = 0$, $u = 0$.</p> <p>When $x = 1$, $u = 1$.</p> $\int_0^1 f\left(\frac{1}{5}x^2 + \frac{4}{5}x\right)(x+2)dx = \int_0^1 f(u)\frac{5}{2}du$ $= \frac{5}{2} \int_0^1 f(u)du$ $= \frac{5}{2} A$
12	C	<p>The period of the product function is a multiple of the periods of the two functions.</p> <p>Period $f(x) = \frac{2\rho}{a}$</p> <p>Period $g(x) = \frac{\rho}{c}$</p> <p>Option A</p> <p>ac cannot be a multiple of the period of $f(x)$ because $ac \cdot \frac{2\rho}{a} = \frac{a^2c}{2\rho}$.</p> <p>$\frac{ac^2}{2\rho}$ cannot be an integer regardless of the values of a and c.</p> <p>Option B</p> <p>Similarly, $2ac$ cannot be a multiple of the period of $f(x)$ because $2ac \cdot \frac{2\rho}{a} = \frac{a^2c}{\rho}$ is not an integer.</p>

		<p>Option C</p> $\left. \begin{array}{l} 2\rho, \frac{2\rho}{a} = a \\ 2\rho, \frac{\rho}{c} = 2c \end{array} \right\} \Rightarrow 2\rho \text{ could be the period of } h(x) \Rightarrow 2\pi \text{ could be the period of } h(x)$ <p>Option D</p> <p>$\frac{2\rho}{ac}$ cannot be a multiple of the period of $f(x)$ because $2ac, \frac{2\rho}{ac} = \frac{a^2c^2}{\rho}$ is not an integer.</p> <p>Option E</p> <p>$\frac{\rho}{ac}$ cannot be a multiple of the period of $f(x)$ because $2ac, \frac{\rho}{ac} = \frac{2a^2c^2}{\rho}$ is not an integer.</p>
13	E	<p>Determine the coordinates of the point of intersection between the two functions, to the right of the y-axis.</p> $f(x) = g(x)$ $e^x - x = 3$ $x = 1.5052415$ <p>The volume generated by the rotation about the x-axis is given by the formula</p> $\begin{aligned} V &= \rho \int_a^b [g(x) - f(x)]^2 dx \\ &= \rho \int_0^{1.505} (3 - e^x + x)^2 dx \\ &= 11.14 \end{aligned}$
14	A	<p>If the particle is in equilibrium, the $\mathbf{F}_1 + \mathbf{F}_2 + \mathbf{F}_3 = 0$</p> $2\mathbf{i} + 3\mathbf{j} - \mathbf{i} - 4\mathbf{j} + \mathbf{F}_3 = 0$ $\mathbf{F}_3 = -\mathbf{i} + \mathbf{j}$ $ \mathbf{F}_3 = \sqrt{2}$
15	B	<p>A confidence interval is given by</p> $\left(\bar{x} - z \frac{s}{\sqrt{n}}, \bar{x} + z \frac{s}{\sqrt{n}} \right)$ $\frac{z}{\sqrt{200}} = 0.1386$ $\begin{aligned} z &= 0.1386 \cdot \sqrt{200} \\ &= 1.96 \end{aligned}$ <p>$z = 1.96$ corresponds to a 95% confidence interval</p>

16	B	$ \begin{aligned} z &= (\sqrt{3} - i)(1 + i)(-2i) \\ &= 2\text{cis}\left(-\frac{\rho}{6}\right) \times \sqrt{2}\text{cis}\left(\frac{\rho}{4}\right) \times 2\text{cis}\left(-\frac{\rho}{2}\right) \\ &= 4\sqrt{2}\text{cis}\left(-\frac{\rho}{6} + \frac{\rho}{4} - \frac{\rho}{2}\right) \\ &= 4\sqrt{2}\text{cis}\left(-\frac{5\rho}{12}\right) \end{aligned} $
17	D	<p>Option A</p> $ \begin{aligned} \mathbf{u} \cdot \mathbf{v} &= (2ai + bj) \cdot (bi + aj) \\ &= 2ab + ab \\ &= 3ab \dots \text{true} \end{aligned} $ <p>Option B</p> <p>Vectors \mathbf{u} and \mathbf{v} are collinear therefore there is at least one real number $k \neq 0$ such that $\mathbf{v} = k\mathbf{u}$.</p> $\mathbf{v} = k\mathbf{u} \Rightarrow 2ai + bj = k(bi + aj)$ <p>Equate corresponding coefficients.</p> $b = 2ka \dots [1]$ $a = kb \dots [2]$ <p>From equation [2], $k = \frac{a}{b}$.</p> <p>Substitute the expression for a into equation [1].</p> $b = 2 \cdot \frac{a}{b} \cdot a \Rightarrow b^2 = 2a^2 \dots \text{true}$ <p>Option C</p> <p>From $b^2 = 2a^2$, $\frac{b^2}{a^2} = 2 \dots \text{true}$</p> $\Rightarrow \frac{b}{a} = \pm\sqrt{2}$ <p>Option D</p> $ \begin{aligned} \mathbf{u} &= \sqrt{(2a)^2 + b^2} \\ &= \sqrt{4a^2 + b^2} \\ \mathbf{v} &= \sqrt{a^2 + b^2} \\ \mathbf{u} &\neq \mathbf{v} \dots \text{incorrect option} \end{aligned} $ <p>Option E</p> $ \mathbf{v} = \sqrt{a^2 + b^2} \dots \text{true}$

18	E	<p>Option A</p> $\begin{aligned} z &= \text{cis}(\theta) = \cos(\theta) + i\sin(\theta) \\ \bar{z} &= \cos(\theta) - i\sin(\theta) \\ &= \cos(-\theta) + i\sin(-\theta) \\ &= \text{cis}(-\theta) \dots \text{true} \end{aligned}$ <p>Option B</p> $\begin{aligned} z^2 &= [\text{cis}(\theta)]^2 \\ &= \text{cis}(2\theta) \dots \text{true} \end{aligned}$ <p>Option C</p> $\begin{aligned} \bar{z} &= \sqrt{\cos^2(\theta) + \sin^2(\theta)} \\ &= 1 \dots \text{true} \end{aligned}$ <p>Option D</p> $\begin{aligned} \frac{z}{\bar{z}} &= \frac{\text{cis}(\theta)}{\text{cis}(-\theta)} \\ &= \text{cis}(\theta + \theta) \\ &= \text{cis}(2\theta) \dots \text{true} \end{aligned}$ <p>Option E</p> $\begin{aligned} z\bar{z} &= \text{cis}(\theta)\text{cis}(-\theta) \\ &= \text{cis}(\theta - \theta) \quad \text{or} \quad z\bar{z} = (z)^2 \\ &= \text{cis}(0) \quad = 1 \\ &= 1 \end{aligned}$
19	B	$a = 2v^2 - v, \text{ where } a = \frac{dv}{dt} = 2v^2 - v$ $\frac{dv}{2v^2 - v} = dt$ <p style="text-align: right;">Using partial fractions,</p> $\begin{aligned} \frac{1}{2v^2 - v} &= \frac{1}{v(2v - 1)} \\ &= \frac{2}{2v - 1} - \frac{1}{v} \end{aligned}$ $\int_1^v \left(\frac{2}{2v - 1} - \frac{1}{v} \right) dv = \int_0^t dt$ $\left[\log_e(2v - 1) - \log_e(v) \right]_1^v = t$ $\log_e \left(\frac{2v - 1}{v} \right) = t$ $\frac{2v - 1}{v} = e^t$ $2v - 1 = ve^t$ $2v - ve^t = 1$ $v = \frac{1}{2 - e^t}$

20	C	$F = ma \Rightarrow a = \frac{16}{m} \text{ m/s}^2$ $s = \frac{1}{2}(u + v)$ $12 = \frac{1}{2}(u + 20)$ $\triangleright u = 4 \text{ m/s}$ $v = u + at$ $20 = 4 + \frac{16}{m} \cdot 4$ $\triangleright m = 4 \text{ kg}$
21	C	<p>The new random variable is $(100\% + 30\%)$ of X plus an extra \$2.</p> $(100 + 30) \% = 130\% = 1.3$ <p>Therefore, the random variable for the new charges is $1.3X + 2$.</p>
22	B	$z_1 = 1 \Rightarrow 1 - a + b = 0$ $\Rightarrow a = 1 + b \dots [1]$ $z_2 = 1 - i \Rightarrow (1 - i)^4 - a(1 - i) + b = 0$ $\Rightarrow -4 - a + ai + b = 0 \dots [2]$ <p>Substitute [1] into [2].</p> $-4 - (1 + b) + (1 + b)i + b = 0$ $-4 - 1 - b + i + bi + b = 0$ $-5 + i + bi = 0$ $bi = 5 - i$ $b = 5i + 1$ $a = 1 + b$ $= 2 + 5i$

SECTION 2**Question 1 (12 marks)****a.**

$$a = \frac{1}{6}, b = \frac{1}{3}$$

1A**b.**

$$E(x) = 1 \cdot \frac{1}{6} + 2 \cdot \frac{1}{3} + 3 \cdot \frac{1}{2}$$

1M

$$E(x) = \frac{7}{3}$$

1A**c.**

$$\text{Var}(X) = E(X^2) - E(X)^2$$

1M

$$\text{Var}(X) = 1^2 \times \frac{1}{6} + 2^2 \times \frac{1}{3} + 3^2 \times \frac{1}{2} - \left(\frac{7}{3}\right)^2$$

1A

$$= \frac{36}{6} - \frac{49}{9}$$

$$= \frac{5}{9}$$

d.

y	2	3	4	5	6
$\Pr(Y = y)$	$\frac{1}{36}$	$\frac{1}{9}$	$\frac{5}{18}$	$\frac{1}{3}$	$\frac{1}{4}$

1A

$$\begin{aligned} \Pr(Y = 2) &= \frac{1}{6} \cdot \frac{1}{6} & \Pr(Y = 4) &= \frac{1}{3} \cdot \frac{1}{3} + \frac{1}{6} \cdot \frac{1}{2} + \frac{1}{2} \cdot \frac{1}{6} \\ &= \frac{1}{36} & &= \frac{5}{18} \end{aligned}$$

1M**e.**

$$\begin{aligned} E(Y) &= 2 \cdot \frac{1}{36} + 3 \cdot \frac{1}{9} + 4 \cdot \frac{5}{18} + 5 \cdot \frac{1}{3} + 6 \cdot \frac{1}{4} \\ &= \frac{14}{3} \end{aligned}$$

1A

$$\begin{aligned} E(Y^2) &= 4 \cdot \frac{1}{36} + 9 \cdot \frac{1}{9} + 16 \cdot \frac{5}{18} + 25 \cdot \frac{1}{3} + 36 \cdot \frac{1}{4} \\ &= \frac{1}{9} + 1 + \frac{40}{9} + \frac{25}{3} + 9 \\ &= \frac{206}{9} \end{aligned}$$

1A

$$\begin{aligned}
 E(Y)^2 + E(Y) - \frac{E(Y^2)}{2} &= \left(\frac{14}{3}\right)^2 + \frac{14}{3} - \frac{1}{2} \times \frac{206}{9} \\
 &= \frac{196}{9} + \frac{14}{3} - \frac{103}{9} \\
 &= \frac{93}{9} + \frac{42}{9} \\
 &= \frac{135}{9} \\
 &= 15 \dots \text{as required}
 \end{aligned}$$

1A**f.**

Median occurs at 0.5.

1M


$$\frac{1}{4} = 0.25$$

$$\frac{1}{3} + \frac{1}{4} = \frac{7}{12} = 0.58$$

The median is 5.

1A

Alternatively, $\frac{1}{36} + \frac{1}{9} + \frac{5}{18} = 0.42$ and $\frac{1}{36} + \frac{1}{9} + \frac{5}{18} + \frac{1}{3} = 0.75$.

Question 2 (12 marks)**a.****1A**The x and y intercepts are both at $(0, 0)$.

Two vertical asymptotes: $x = -\frac{\rho}{2}$ and $x = \frac{\rho}{2}$

1A**b.**

$$f(x) = \tan(x)e^{2x}$$

$$\begin{aligned}
 f'(x) &= \frac{1}{\cos^2(x)} e^{2x} + \tan(x) \times 2e^{2x} \\
 &= \frac{1}{\cos^2(x)} e^{2x} + 2\tan(x)e^{2x} \\
 &= \frac{1}{\cos^2(x)} e^{2x} + 2f(x)
 \end{aligned}$$

1A

$$f'''(x) = -2\cos^{-3}(x)(-\sin(x))e^{2x} + \frac{1}{\cos^2(x)} \times 2e^{2x} + 2f'(x) \quad \text{1A}$$

$$f''(x) = 2\tan(x)\sec^2(x)e^{2x} + 2\sec^2(x)e^{2x} + 2f'(x)$$

$$f''(x) = 2f'(x) + 2\sec^2(x)[f(x) + e^{2x}] \dots \text{as required}$$

1A

c.

Points of inflection occur when $\frac{d^2y}{dx^2} = 0$. 1A

The point of inflection has coordinates $(-0.785, -0.208)$.

To one decimal place, $(-0.8, -0.2)$.

1A

d.

When $x = \frac{\rho}{4}$, $y = f\left(\frac{\rho}{4}\right)$ 1A

$$= \tan\left(\frac{\rho}{4}\right) e^{2 \times \frac{\rho}{4}}$$

$$= e^{\frac{\rho}{2}}$$

$$\left. \frac{dy}{dx} \right|_{x=\frac{\rho}{4}} = 4e^{\frac{\rho}{2}} \quad \text{1A}$$

e.

The equation of the tangent at $x = \frac{\rho}{4}$ is

$$y - e^{\frac{\rho}{2}} = 4e^{\frac{\rho}{2}} \left(x - \frac{\rho}{4} \right)$$

$$y = e^{\frac{\rho}{2}} + 4e^{\frac{\rho}{2}}x - \frac{\rho}{4}4e^{\frac{\rho}{2}}$$

$$y = 4e^{\frac{\rho}{2}}x + e^{\frac{\rho}{2}}(1 - \rho)$$

$$m = 4e^{\frac{\rho}{2}}, c = e^{\frac{\rho}{2}}(1 - \rho)$$

1M

2A

Question 3 (12 marks)

a.


$$\begin{aligned} |z + 4i - 1| &= |x + iy + 4i - 1| \\ &= \sqrt{(x - 1)^2 + (y + 4)^2} \end{aligned} \quad \text{1M}$$

$$\sqrt{(x - 1)^2 + (y + 4)^2} = 3 \dots \text{square both sides of the equation}$$

$$(x - 1)^2 + (y + 4)^2 = 9$$

The set of complex numbers P represents a circle with radius 3 and centre $(1, -4)$.

1A

b.**1A + 1A**

The shaded region satisfies the conditions given including the boundary of the circle.

c.

The solutions of the equation $|z + 2 + 4i| = |z - 4 + 4i|$ represent the locus of points at the same distance from $-2 - 4i$ and $4 - 4i$.

The locus of points is the median bisector of the line segment passing through $(-2, -4)$ and $(4, -4)$ with equation $x = 1$.



1A

The points needed are in fact the solutions of the system of equations

$$\begin{cases} x = 1 \\ (x - 1)^2 + (y + 4)^2 = 9 \end{cases} \Rightarrow \begin{cases} x = 1, y = -7 \\ x = 1, y = -1 \end{cases}$$

$$z_1 = 1 - 7i$$

$$z_2 = 1 - i$$

**1A****d.****1A**

e. i.

Solve the system of simultaneous equations

$$\begin{cases} (x - 1)^2 + (y + 4)^2 = 9 \dots [1] \\ (x - 0)^2 + (y + 4)^2 = c \dots [2] \end{cases} \dots \text{subtract equation [2] from [1]} \quad \mathbf{1M}$$

$$(x - 1)^2 - x^2 = 9 - c$$

$$x^2 - 2x + 1 - x^2 = 9 - c$$

$$-2x = 8 - c \quad \mathbf{1A}$$

$$x = \frac{c}{2} - 4$$

Substitute the value of x into equation [2].

$$\left(\frac{c}{2} - 4\right)^2 + (y + 4)^2 = c$$

$$(y + 4)^2 = c - \frac{c^2}{4} + 4c - 16$$

$$(y + 4)^2 = 5c - \frac{c^2}{4} - 16$$


$$y + 4 = \pm \sqrt{5c - \frac{c^2}{4} - 16}$$

$$y = -4 \pm \sqrt{5c - \frac{c^2}{4} - 16}$$

The second circle has to be inside the first circle. Therefore, $y = -4 + \sqrt{5c - \frac{c^2}{4} - 16}$ is the expression that satisfies this condition. $\mathbf{1A}$

e. ii.For the largest value of c , the two circles must only have one point in common. The centre of the inside circle is $(0, -4)$, therefore its maximum radius is 2.

$$c = 2^2 = 4. \quad \mathbf{1A}$$

e. iii.

Question 4 (9 marks)**a.**

The object moves under constant acceleration.

$$v = u + at, \text{ where } u = 0, t = 10 \text{ s and } a = 3 \text{ m/s}^2.$$


$$v = 0 + 3 \times 10 = 30 \text{ m/s}$$

1A**b.**

$$s = \frac{1}{2}(u + v)t$$

$$= \frac{1}{2} \cdot 30 \cdot 10$$

$$= 150 \text{ m}$$

1A**c.****2A****d.**

The object is moving under constant deceleration (negative acceleration) to come to a stop.

$u = 30\cos(12^\circ)$ is the velocity at the end of the horizontal surface and the horizontal component at the beginning of the inclined plane.

$$v = 0 \text{ (at the end of the motion on the inclined plane)}$$

$$v^2 = u^2 + 2as$$

$$0 = [30\cos(12^\circ)]^2 + 2a \times 25$$

$$a = -17.22 \text{ ms}^{-2}$$

1M + 1A**e.**

$$\text{The horizontal components: } 2a = F\cos(12^\circ) + 2g\cos(78^\circ) - F_{fr} \quad \text{1A}$$

$$\text{The vertical components: } N + F\sin(12^\circ) = 2g\cos(12^\circ) \quad \text{1A}$$

$$\begin{cases} 2a = F\cos(12^\circ) + 2g\cos(78^\circ) - F_{fr} \\ N + F\sin(12^\circ) = 2g\cos(12^\circ) \end{cases}$$

$$\begin{cases} 2 \times (-17.22) = 6\cos(12^\circ) + 2g\cos(78^\circ) - m(2g\cos(12^\circ) - 6\sin(12^\circ)) \\ N = 2g\cos(12^\circ) - 6\sin(12^\circ) \end{cases}$$

$$m = \frac{-2 \cdot (-17.22) + 6\cos(12^\circ) + 2g\cos(78^\circ)}{2g\cos(12^\circ) - 6\sin(12^\circ)} \quad \text{1A}$$

$$= 2.48$$

Question 5 (13 marks)**a.**At $t = 0$, $\mathbf{r}(0) = \mathbf{0}$.

$$\begin{cases} 3\sin(0) - \sqrt{3}\cos(0) + a = 0 \\ -\sqrt{3}\sin(0) - 3\cos(0) + b = 0 \end{cases}$$

1M

$$\begin{cases} -\sqrt{3} + a = 0 \\ -3 + b = 0 \end{cases}$$

$$a = \sqrt{3} \text{ and } b = 3$$

1A**b.**

$$\begin{aligned} |\mathbf{r}(t)| &= \left| \left[3\sin(t) - \sqrt{3}\cos(t) + \sqrt{3} \right] \mathbf{i} + \left[-\sqrt{3}\sin(t) - 3\cos(t) + 3 \right] \mathbf{j} \right| \\ &= \sqrt{\left[3\sin(t) - \sqrt{3}\cos(t) + \sqrt{3} \right]^2 + \left[-\sqrt{3}\sin(t) - 3\cos(t) + 3 \right]^2} \\ &= \sqrt{-24\cos(t) + 14} \end{aligned}$$

1M

$$\sqrt{-24\cos(t) + 14} = 6$$

$$-24\cos(t) + 14 = 36$$

$$-24\cos(t) = 12$$

$$\cos(t) = -\frac{1}{2} \Rightarrow t = \left\{ \frac{2\rho}{3}, \frac{4\rho}{3} \right\}$$

$$t_1 = 2.09 \text{ s and } t_2 = 4.19 \text{ s}$$

1A**c.**

$$\mathbf{r}(t) = \left[3\sin(t) - \sqrt{3}\cos(t) + \sqrt{3} \right] \mathbf{i} + \left[-\sqrt{3}\sin(t) - 3\cos(t) + 3 \right] \mathbf{j}$$

$$\dot{\mathbf{r}}(t) = \left[3\cos(t) + \sqrt{3}\sin(t) \right] \mathbf{i} + \left[-\sqrt{3}\cos(t) + 3\sin(t) \right] \mathbf{j}$$

1A

$$\ddot{\mathbf{r}}(t) = \left[-3\sin(t) + \sqrt{3}\cos(t) \right] \mathbf{i} + \left[\sqrt{3}\sin(t) + 3\cos(t) \right] \mathbf{j}$$

$$\begin{aligned} |\ddot{\mathbf{r}}(t)| &= \sqrt{\left[-3\sin(t) + \sqrt{3}\cos(t) \right]^2 + \left[\sqrt{3}\sin(t) + 3\cos(t) \right]^2} \\ &= \sqrt{12\sin^2(t) + 12\cos^2(t)} \\ &= \sqrt{12} \dots \text{as required} \end{aligned}$$

1A**d.**

$$\dot{\mathbf{r}}(t) = \left[3\cos(t) + \sqrt{3}\sin(t) \right] \mathbf{i} + \left[-\sqrt{3}\cos(t) + 3\sin(t) \right] \mathbf{j}$$

$$m = \sqrt{3^2 + \sqrt{3}^2}$$

$$= \sqrt{12}$$

$$= 2\sqrt{3}$$

1A

$$\alpha = \tan^{-1}\left(\frac{\sqrt{3}}{3}\right) \text{ and } b = \tan^{-1}\left(-\frac{3}{\sqrt{3}}\right)$$

$$= 0.52 \qquad \qquad = -1.05$$

1A

$$\dot{r}(t) = \left[2\sqrt{3} \cos(t + 0.52) \right] \mathbf{i} - \left[2\sqrt{3} \cos(t - 1.05) \right] \mathbf{j}$$

1A**e.**

$$\begin{cases} x = 3\sin(t) - \sqrt{3}\cos(t) + \sqrt{3} \\ y = -\sqrt{3}\sin(t) - 3\cos(t) + 3 \end{cases}$$

$$\begin{cases} x - \sqrt{3} = 3\sin(t) - \sqrt{3}\cos(t) \dots \text{multiply by } \sqrt{3} \\ y - 3 = -\sqrt{3}\sin(t) - 3\cos(t) \end{cases}$$

$$\begin{cases} \sqrt{3}x - 3 = 3\sqrt{3}\sin(t) - 3\cos(t) \dots [1] \\ y - 3 = -\sqrt{3}\sin(t) - 3\cos(t) \dots [2] \end{cases} \dots \text{subtract [1] - [2]}$$

$$\sqrt{3}x - 3 - y + 3 = 3\sqrt{3}\sin(t) + \sqrt{3}\sin(t)$$

$$\sin(t) = \frac{\sqrt{3}x - y}{4\sqrt{3}} \dots \text{rationalise the denominator}$$

1M

$$\sin(t) = \frac{3x - \sqrt{3}y}{12} \dots [3]$$

1A

Substitute [3] in [2].

$$y - 3 = -\sqrt{3} \cdot \frac{3x - \sqrt{3}y}{12} - 3\cos(t)$$

$$3\cos(t) = 3 - y - \sqrt{3} \cdot \frac{3x - \sqrt{3}y}{12}$$

$$\cos(t) = 1 - \frac{y}{3} - \frac{\sqrt{3}x - y}{12}$$

$$\cos(t) = \frac{12 - 4y - \sqrt{3}x + y}{12}$$

$$\cos(t) = \frac{12 - 3y - \sqrt{3}x}{12} \dots [4]$$

1A

$$\sin^2(t) + \cos^2(t) = 1$$

$$\left(\frac{3x - \sqrt{3}y}{12}\right)^2 + \left(\frac{12 - 3y - \sqrt{3}x}{12}\right)^2 = 1$$

$$12x^2 + 12y^2 - 12\sqrt{3}xy - 72y - 24\sqrt{3}x = 0$$

$$x^2 + y^2 - \sqrt{3}xy - 6y - 2\sqrt{3}x = 0$$

1A